

CEBI WORKING PAPER SERIES

Working Paper 09/20

THE TIMING OF EARLY INTERVENTIONS AND
CHILD AND MATERNAL HEALTH

Jonas Lau-Jensen Hirani
Hans Henrik Sivertsen
Miriam Wüst

ISSN 2596-44TX

CEBI

Department of Economics
University of Copenhagen
www.cebi.ku.dk

The Timing of Early Interventions and Child and Maternal Health

Jonas Lau-Jensen Hirani*, Hans Henrik Sievertsen[†] and Miriam Wüst[‡]

Version: March 23, 2020.

Abstract

What is the impact of timing of early-life investment policies on child and maternal health? Exploiting variation from a 2008 Danish nurse strike, we study this question in the context of universal nurse home visiting. We show that early but not later strike exposure increases child (and mother) general practitioner contacts in the first four years after childbirth. Mothers, who forgo an early nurse visit (rather than a later one), have a higher probability of mental health specialist contacts in the first two years after childbirth. We highlight two channels for these results, screening and information provision: We show that nurses perform well in identifying maternal mental health risks during early home visits in control years (likely preventing longer-term problems). Finally, we show that first-born children and children of parents without a health-related education drive our results. A stylized calculation confirms that short-run health benefits from early universal home visiting outweigh costs.

JEL Codes: I11, I12, I14, I18, I21

Keywords: Early-life health, early interventions, nurse home visiting, parental investments

*Department of Economics, University of Copenhagen and The Danish Center for Social Science Research (VIVE). Herluf Trolles Gade 11, 1052 Copenhagen, email: jjh@vive.dk.

[†]Department of Economics, University of Bristol and The Danish Center for Social Science Research (VIVE). Priory Road Complex, Priory Road, Bristol BS8 1TU, UK, email: h.h.sievertsen@bristol.ac.uk.

[‡]Corresponding author: University of Copenhagen, Center for Economic Behavior and Inequality (CEBI), Department of Economics and The Danish Center for Social Science Research (VIVE). Øster Farismagsgade 5, 1353 Copenhagen, email: miriam.w@econ.ku.dk., phone: +45 35321365

We gratefully acknowledge financial support from the Innovation Foundation Denmark grant 5155-00001B. The Copenhagen City Archives generously provided the data on nurse home visiting in the municipality of Copenhagen. We thank Boel Leth Emanuel (municipality of Copenhagen) for valuable input on the data. We thank Søren Leth Pedersen, Maya Rossin-Slater, Christine Valente, Stephanie Vincent Lyk-Jensen and seminar participants at briq (Bonn), SOFI (Stockholm), CEBI (University of Copenhagen), Copenhagen Business School, the NBER Children's Program Meeting 2019, the Danish Ministry of Health, CINCH Essen and at the 2019 EALE conference for helpful comments.

1 Introduction

This paper studies the importance of the timing of early-life public health investments for child and maternal health. Evidence on this question is sparse but instrumental for policy: In the light of mounting evidence on the importance of early interventions for the short- and long-run health and human capital of children, policy decisions in many settings evolve around central features of early-life investment program, such as their timing.

We study an investment program in Denmark, where all new families are eligible for up to five universal home visits by a trained nurse during the first year of a child’s life. These visits focus on health screening, the provision of information, and counseling to new parents (on topics such as infant feeding, infant development and child-parent interactions). Additionally, nurses refer families with identified needs to other health care professionals, such as general practitioners (GPs) or hospitals.

To identify the impact of timing of nurse home visiting (NHV), we exploit exposure to a 2008 nurse strike for families with children born in the seven months *prior* to the strike. During the 61 days of the strike, the vast majority of non-emergency nurse care was cancelled. Importantly, cancelled visits were not rescheduled. We exploit the strike-induced variation in nurse home visits, together with information on children born in non-strike years in a difference-in-differences design.

To make our study feasible, we collected individual-level data on program take-up (number and timing of nurse visits) in largest municipality in Denmark (Copenhagen) and linked these records to administrative data on family background and health outcomes.¹ Thus we break new grounds by compiling data on actual program take-up (the policy-relevant margin in a universal program), which allows us to be specific about the intensity of the treatment that we study. The link to administrative data allows us to analyze the credibility of our empirical design (by assessing the compliance with the nurse strike across different groups of families).

¹While Scandinavia is well-known for its high-quality administrative data in many domains, national administrative data sources typically lack individual-level data on municipal programs—such as NHV, nurseries or preschools.

In our first set of results, we show that, while children born in the 210 days prior to the strike on average missed one scheduled postnatal nurse visit, depending on their date of birth *relative* to the strike, children had a different age at the forgone visit. In Copenhagen, nurses only performed ten percent of the home visits that we observe in the same weeks in the years prior to and after the strike. Exploiting the merged nurse records and administrative data on family background, we show that the strike impacted families similarly across characteristics, likely observed by nurses. This finding illustrates the broad coverage of the strike in the population and relieves concerns that nurses to a large degree chose the families that would forgo their visit. Additionally, we show that (given that all children were born before the onset of the strike) other aspects of care around birth (such as prenatal midwife contacts or hospital admissions at birth) were not impacted by children’s strike exposure.

Moving on to our reduced form analysis of the impact of strike exposure at specific ages, we show that exposure during the initial months of a child’s life is relatively more influential for child and maternal health than later exposure. We measure health by the uptake of additional medical care: Children, who were born in the two to three months up to the strike, and thus were likely to miss the early nurse visits due to the strike, have more contacts to general practitioners (GP) in the first four years of life than children, who were older at their exposure to the strike. This result holds for both regular and emergency GP contacts (the latter not being performed by the family’s regular GP and outside GP office hours). Moreover, our results for yearly measures of GP contacts confirm this finding, i.e. our results are not driven by a closer relationship with the family GP or a substitution of nurse visits with GP visits during the strike period.² We also find suggestive evidence for a higher probability of hospital contacts in the second year of the child’s life for early strike-exposed children. This finding further substantiates that our results for an increased uptake of health care reflect children’s underlying health.

We also study maternal health care usage as a consequence of strike exposure. First,

²Our main outcome measures of GP contacts exclude preventive care contacts to the GP, which we study separately.

we find that mothers, who are likely to forgo an early nurse visit due to the strike, have more GP contacts in the first four years after their child’s birth. Second, we find suggestive evidence for early strike-exposed mothers being *more* likely to have at least one contact with a psychologist or psychiatrist in the first two years of the child’s life. This finding suggests that early strike-exposed mothers (who thus lack a nurse visit focusing on early screening for mental health issues) end up receiving more specialist treatment. While missing an early nurse visit initially (and mechanically) may result in fewer mothers being referred to other specialists, our findings suggest that (in the longer run) early strike exposure leads to an increased likelihood of mothers experiencing mental health problems that require specialist attention. Thus our finding is in line with recent studies documenting the importance of different aspects of the early home environment (in our case the early detection and prevention of severe problems) for maternal postpartum mental health (Butikofer et al., 2018; Baranov et al., 2019; Persson and Rossin-Slater, 2019).

Having established health effects of missing an early nurse visit, we consider two possible mechanisms by exploring the role of screening and information. First, early nurse visits may help to identify adverse conditions in a timely fashion and prompt additional treatments. We use data from the non-strike exposed control period to show that, at initial visits, nurses typically record issues related to feeding, child physical health and maternal well-being. In control year data, these initial registrations correlate with both future nurse registrations of health issues and the increased use of health care services among children, as well as the likelihood of future maternal psychiatric contacts. These correlations suggest that early nurse visits act as an important screening device to identify vulnerable children and mothers. In absence of early nurse visits, for the marginal child and mother, health problems may go unnoticed for a longer period and contribute to longer-term adverse health effects. Our results for the impact of early strike-exposure on maternal contacts to psychologists or psychiatrists are in line with this reasoning. Moreover, given documented correlations of maternal postnatal mental health and child-parent interactions and child development (Cooper and Murray,

1998; Lovejoy et al., 2000; Paulson et al., 2006; Wachs et al., 2009), screening for postnatal maternal well-being issues may be one driver also for impacts of early NHV on children.

Second, in the absence of early nurse visits, parents may lack specific information, which is typically provided by nurses and is difficult to replace by other and less specialized health care providers, such as GPs. Moreover, information and counseling provided by nurses may impact parents' investment behaviors, such as breastfeeding, parent-child interactions or uptake of other preventive care. To examine the relevance of this channel, we study the impact of strike exposure among children across different backgrounds. We find suggestive evidence that higher parity children and children of parents with an educational background in a health-related field (nurses, midwives, doctors and pedagogues) are less affected by strike exposure than their first-parity and not health-educated counterparts, respectively. At the same time, we find no strong and unambiguous evidence for a socio-economic gradient in the effect of early strike exposure. These findings indicate that at least part of the beneficial effect of early NHV runs through a specific information channel. While we study parents' participation in the vaccination and preventive care programs (as our main measures of parental investment behaviors), we do not detect a strong impact of the timing of nurse visits in our design. However, these analyses are constrained by power issues.

Our stylized analysis of the direct costs and benefits of early nurse visits relative to later visits (based on a limited set of outcomes from the domain of health and thus ignoring other potential benefits of earlier nurse visits) shows that immediate benefits (in terms of averted child and mother GP visits) clearly outweigh costs (with 331-426 EUR). Thus our findings indicate that early universal visits are a cost-effective intervention to promote children's and mothers' health in settings that resemble the Danish health care system. Given our findings, universal child programs should have a strong focus on the initial period of family formation after the birth of a child.

Our work contributes to a large literature documenting causal links between childhood experiences—shocks and exposure to policies—and later life outcomes (for an overview see

Almond and Currie, 2011; Almond et al., 2018). We make three contributions: First, when studying the causal effects of early-life investment programs (such as nurse home visiting, or childcare and early education provisions), the majority of work has considered the effects of program exposure. However, we still lack insights on the causal effects of important design aspects of early-life investment policies, such as timing or intensity. In our paper, rather than studying the margin of program exposure, we consider the so far largely unexplored impacts of *the relative timing* of access to early-life health programs. Our study extends earlier work by Kronborg et al. (2016), who study mothers giving birth *during* and shortly prior to the nurse strike and only find short-lived effects of strike exposure on the take-up of GP care for children. However, in their paper all strike-exposed mothers and children forgo the earliest home visits (the ones that we show are influential) and vary in their access to various treatments: prenatal midwife consultations, hospital stays after birth and the early postnatal nurse visits.³

Second, a large share of the work on early-life investment policies has been set in a U.S. context and as a consequence has considered *targeted* programs.⁴ Existing work on NHV has primarily focused on contemporary targeted programs as well (Olds et al., 1986, 1998, 2002; Vaithianathan et al., 2016; Doyle et al., 2015; Sandner et al., 2018; Sandner, 2019).⁵

³Strike-exposed mothers in their analysis received less pre- and postnatal care: Mothers, who gave birth during the strike received fewer prenatal midwife consultations, were more likely to be discharged from hospital on the day of birth, and received fewer nurse home visits. Mothers, who gave birth in the two weeks prior to the strike had higher probability of not receiving the initial nurse visit but were unaffected with respect to the access of prenatal care. Mothers, who gave birth earlier (two weeks to two months prior to the strike) were unaffected with respect to the prenatal care offers, hospital care around birth but had an increased probability of a canceled second nurse visit. Given that all mothers in the sample lack the early home visits after birth, our analysis identifies a different margin of treatment (focusing only on the importance of timing of postnatal care). Moreover, while Kronborg et al. (2016) cannot link data on NHV to data on family background and health outcomes, we perform a complier analysis, i.e. assess the “coverage” of strike exposure in the population of families. Finally, we both analyze a broader set of relevant outcomes (including maternal well-being) and the potential channels for our main results.

⁴Examples include RCT studies on the targeted Perry Preschool Program, the Abecedarian project (Heckman et al., 2013; Conti et al., 2016), and observational studies on the short- and long-run impact of Head Start (Currie and Thomas, 1995; Garces et al., 2002; Masse and Barnett, 2002; Schweinhart et al., 2005; Belfield et al., 2006; Ludwig and Miller, 2007; Anderson, 2008; Deming, 2009; Heckman et al., 2010a,b; Carneiro and Ginja, 2014; Campbell et al., 2014; García et al., 2016; De Haan and Leuven, 2016; García et al., 2017; Thompson, 2017). Also in a US context, there are a few examples for studies considering universal provision of preschool (see, for example, Cascio, 2009, 2015).

⁵Existing evidence suggests that targeted NHV can be effective in improving a large range of short- and

However, many countries offer *universal* programs and the results from studies on targeted programs do not easily generalize to settings with universal implementation. Our study is the first to analyze the causal impacts of a contemporary universal program.⁶ Evidence on the impact of universal health programs and their design is instrumental for policy design in many settings.

Third, we shed light on two relevant mechanisms for the impact of timing of NHV on child and mother health: Screening (and potential referral of families to other health professionals) and information (e.g., about infant feeding or age-specific child-parent interactions). This information may matter in its own right (i.e., it may be new to parents) or modify parental beliefs (i.e., it may help parents to update their reading of information that they have access to). Recent research documents the importance of parental beliefs—their interpretation of rather than their pure awareness of information—for both child health outcomes and parental investment behaviors (see, for example, Cunha et al., 2013; Attanasio et al., 2015; Boneva and Rauh, 2018; Biroli et al., 2018). Our unique data allow us to explore the question of which elements matter in NHV by using specific nurse registrations and the heterogeneity of effects of NHV across different types of parents. While we cannot formally distinguish whether nurses provide new knowledge to parents or modify their beliefs about its importance, we

long-run child outcomes and points to the role of the structure of the programs and the qualifications of service providers (for an overview on existing studies and a discussion of the impact of provider quality, target group and program features, see Almond and Currie, 2011): Focusing on the targeted Nurse Home Visiting Partnership program in the US, Olds et al. (1986, 1998, 2002) show that high-frequency pre- and postnatal visits for at-risk mothers conducted by trained nurses reduced child abuse, decreased children’s emergency room visits and their criminal convictions in adolescence. Similarly, Vaithianathan et al. (2016) provide evidence from New Zealand showing that targeted nurse visits reduced infant mortality and increased both vaccination rates and children’s participation in early childhood education. Doyle et al. (2015) study the targeted Preparing for Life-program in Ireland and find some positive effects on child health (such as asthma issues) and accidents. Sandner et al. (2018) and Sandner (2019) document that the German implementation of the “Pro Kind” program, a home visiting program for low-income first-time mothers, did not impact child health but had impacts on mothers in the RCT: treated mothers reported lower levels of depression. In the longer run, the program increased fertility and decreased maternal labor supply. Work from developing country contexts highlights the important role for child development and long-run outcomes that intensive home visiting can play (Attanasio et al., 2014; Gertler et al., 2014).

⁶Another line of research has documented positive long-run impacts of the historical introduction of universal NHV in Scandinavia of the 1930s and 40s (Wüst, 2012; Hjort et al., 2017; Bhalotra et al., 2017; Bütikofer et al., 2019). All existing evidence on the causal short- and long-run effects of NHV in Scandinavia comes from historical data and considers the extensive margin of treatment exposure. These studies have documented positive long-run effects on the health and socio-economic outcomes of exposed cohorts.

find that the timely provision of both components of NHV (information, screening) most likely are important for the impacts of the program.

The paper proceeds as follows: Section 2 provides information on the institutional background, the 2008 nurse strike and the data sources that we use. Section 3 presents our empirical strategy and discusses the identifying assumptions. Section 4 presents descriptive and main results and examines their robustness and heterogeneity. Section 5 performs a simple cost-effectiveness analysis. Finally, section 6 concludes.

2 Background and Data

2.1 Institutional Background: Pre and postnatal care in Denmark

In Denmark, pre- and postnatal care is provided in the public health care system and all residents have access to care free of charge. Midwives and general practitioners provide prenatal care that consists of regular consultations during pregnancy.⁷ The majority of uncomplicated births are midwife-assisted and take place in public hospitals only. Hospital births account for around 98 percent of all births.

After hospital discharge, the 98 municipalities are responsible for providing postnatal care in the NHV program. While there is some variation in municipal service levels, the Danish National Board of Health (DNBH) issues guidelines and regulations regarding the number, timing and content of nurse visits. As such, NHV consists of a basic package of services offered to all families with a newborn. Additionally, municipalities can choose to offer supplementary services targeted at specific populations of mothers and children. Those services include additional home visits or other services.⁸ Moreover, Danish GPs provide the child preventive health program and administer recommended vaccines in the vaccination program. The Danish preventive care schedule offers eight (voluntary) GP health checks for

⁷The universal offer consists of 4-7 midwife consultations, 3 GP consultations and 2 ultrasound scans Sundhedsstyrelsen (2007). At-risk pregnancies receive additional care.

⁸These services can include offers such as group interventions, interventions targeted at young parents or parents with specific health issues, or interventions specifically directed at fathers.

all children: at around five weeks, at around five months, and yearly for children aged one through six years (Sundhedsstyrelsen, 2007). Additionally, GPs offer one postpartum health check for mothers. In the first year of the child’s life, the Danish vaccination program for children consists of three rounds, at three, five and twelve months, respectively.⁹

2.2 NHV in Copenhagen

Our study focuses on NHV in Copenhagen, the largest municipality in Denmark with around 500,000 inhabitants and around 8-10,000 yearly live births. Appendix Table A1 presents the main features of NHV in Copenhagen. The default number of universally-offered visits in the program is four: an initial visit shortly after birth (A visit), a two month visit (B visit), a four month visit and an eight month visit (C and D visit). Infants, who are discharged after short hospital stays can receive two A visits.¹⁰ Moreover, nurses can provide additional targeted visits to children and families with identified needs at their discretion. The timing of these additional visits is flexible. Finally, the municipality offers optional visits that are available on the request of parents (visits at ages 1.5 and three years).

Home visits usually last between 30 minutes and one hour. During the visits, nurses provide information and counseling to parents and examine the infant. The visits take their point of departure from a general set of main topics (which are of different importance at different ages of the child) outlined in the national guidelines for NHV. At the same time, those guidelines explicitly state that nurses should focus on the needs of the specific family. Thus nurses have large discretion to focus their time in the family home on what they regard as most important. While some topics, typically related to screening (such as tests for certain infant reflexes, monitoring of maternal postnatal well-being and the monitoring of

⁹Each round consists of two separate vaccinations. First, a combined vaccination to immunize against diphtheria, tetanus, pertussis, polio and hib infection. Second, a pneumococcus bacteria vaccination to prevent infant meningitis. We focus in this paper on the vaccinations given in the first year of life but the vaccination program continues with a number of other vaccinations throughout childhood and adolescence.

¹⁰Especially for higher parity births, discharge on the day of birth is not unusual in Denmark: Among uncomplicated births in our sample, 58 percent of mothers are discharged with their infant on the day of birth.

child weight and height), are part of visits to all families, other topics are only covered if the family or the nurse find them relevant.

Given the variation in families’ needs, nurse registrations are of similar variability: Table 1 illustrates the main topics that structure the universal nurse visits in the child’s first year of life (A-D visits in Copenhagen) and which registrations nurses can make. Importantly, domains that are covered in each visit such as infant feeding have age-specific items that nurses can make registrations on (such as “issues with establishment of breastfeeding” or “issues with the introduction of solid food”).

To illustrate the typical content of the nurse visits, Figure 1 presents nurse registrations made by Copenhagen nurses during or shortly after their home visits to families from our control cohort.¹¹

We aggregate nurse registrations into broader categories and plot for each of those categories the share of families with a recorded issue by type of visit (conditional on having received the visit). As the figure illustrates, the visits focus on different domains: While the share of families with “registrations of an issue in any domain” remains rather stable over the course of the four visits, there are important differences especially when comparing the first two and final two universal visits. During the initial visits, nurses typically record issues related to maternal mental well-being and infant feeding issues. The former is very well-defined, mother-specific and highly correlated for women across visits. The latter is child-related but rather unspecific in its content. While registrations on feeding issues are common during the initial visits, nurse observations and registrations on child developmental problems (a summary measure of various dimensions of child development) are more prevalent in the visits at around four and eight months. Using this characterization of the composite nurse treatment, we will return to the importance of different aspects of the treatment in section 4.4.

¹¹As we will detail in section 2.3, we use data on several cohorts of children and mothers, one of them exposed to the nurse strike. In Figure 1, we focus on non-strike exposed children and mothers as strike-exposed families naturally lack nurse registrations.

Table 1 Overview on main topics at nurse visits and options for nurse registrations in the municipality of Copenhagen.

Topic	Examples for items (some visit-specific)	--- Visit ---			
		A 0-14days	B 2months	C 4 months	D 8 months
Background	Issues related to pregnancy and birth, health risks (parental smoking, alcohol, BMI), family structure, etc	✓			
(1) Postpartum maternal health	Physical and mental well-being, formal depression screening	✓	✓	✓	✓
(2) Feeding	Breastfeeding, supplementary feeding, introduction of solid food	✓	✓	✓	✓
(3) Parent-child interactions	Activities, parental recognition of infant needs/signals	✓	✓	✓	✓
(4) Child signals and reactions	Sleep patterns, mood, smile/contact, differentiating btw adults	✓	✓	✓	✓
(5) Child Examinations					
a. Physical health	Weight and height, jaundice	✓	✓	✓	✓
b. Reflexes	Sucking, crawling, Babinski	✓	✓	✓	✓
c. Tactile sense		✓	✓	✓	✓
d. Head	Size, symmetry	✓	✓	✓	✓
e. Skin and navel	Eczema, color and dryness	✓	✓	✓	✓
f. Gross motor dev.	Infant: holds head, changes from stomach to back, sits alone, attempts to crawl		✓	✓	✓
g. Eye-hand coordination	Infant: puts hand in mouth, sees her own hand, pinch grip		✓	✓	✓
h. Vision	Infant: holds eye contact, follows objects	✓	✓	✓	✓
i. Communication	Infant: smiles, chatters		✓	✓	✓
j. Congenital malformations	Ears, hips, genitals, mouth	✓	✓	✓	✓

Notes: The table illustrates the schedule for topics at the given home visit and the nurses' options for registrations in the registration system used by nurses in Copenhagen in the period that we study. Visits A-D refer to the type of the four scheduled universal visits. Additionally, nurses can offer a targeted pregnancy visit (around week 30 of the pregnancy), visits based on identified needs in the family, and a visit at age 1.5 and 3 years (on parental demand), respectively.




Fig. 1 Share of mothers/children with nurse registrations of issues at a given nurse visit (initial visit through eight months visit)

Notes: The share of children with registered issues in each domain for all children with a performed visit and born between September 17, 2008 - April 15, 2009 (the control period). Each domain aggregates a set of binary measures. Children’s issue indicator is one if at least one binary measure is registered as problematic by the family’s nurse.

2.3 Data and Variable Construction

In our analysis, we use data from two sources. First, we access archived records on the universe of home visits from the municipality of Copenhagen for the 2007-2009 period.¹² These registrations were either completed at the family home (using a laptop) or at the nurse’s office directly after a completed visit. For each visit, the data contain the date and type of visit. Additionally, nurses register their observations regarding factors such as child and maternal health, feeding problems, or relevant risk factors in the family (see Table 1 for

¹²These data come from an archive version of the municipality’s administrative system. The full archive of nurse records from Copenhagen includes data on all visits and examinations of children resident in the municipality from January 1, 2007 to December 31, 2010—a total of 35,213 children. These records were transferred to the Copenhagen city archive due to a change of the software used by the Copenhagen nurses. As we are interested in studying the impact of timing of nurse visits in the first year of the child’s life, we do not consider data from the 2010 cohort as they are right-censored, i.e. we do not observe information on all visits before the end of the data period.

examples of focus areas and registration options at different visits).

Second, using children’s unique social security number, we merge the nurse records with population administrative data from Statistics Denmark for the birth cohorts 2007-2010.¹³ The administrative data contains a large set of parental background characteristics such as educational attainment, income, age, civil status and family links irrespective of co-residence, and municipality of residence and birth records. Moreover, the administrative birth records provide information on measures such as children’s birth weight and length, gestation age, the five minute APGAR-score, hospital of birth identifiers and take-up and number of prenatal midwife contacts.

Using data for the years 2007-2014, we create three sets of health outcome measures from the administrative data: First, to study child and maternal health, we examine both the yearly and accumulated number of GP contacts from child age zero to four. GP contacts include both physical meetings and phone and e-mail correspondence with a GP.¹⁴ Given that we only measure health care usage in our data, we are concerned as to whether we pick up actual impacts of strike exposure on child health: Parents may behave more cautious and—in the short run—substitute nurse care with GP care. In the longer run, parents may continue to demand more care, for example, because they build a strong relationship with their family GP due to increased initial contacts.

While we cannot fully disentangle true health effects from alternative explanations for changes in health care take-up, we attempt to provide more insights by dividing our measure of GP contacts into two categories: i) regular (scheduled) GP contacts that typically involve the family GP, and ii) emergency GP contacts (i.e., GP contacts on weekends or outside default opening hours, which are not performed by the family GP).¹⁵ While not perfectly

¹³In our reduced form analysis of strike exposure on child outcomes, we use an additional cohort of children (2010) in our control group. Our results are not sensitive to the choice of control years, as detailed in section 4.5.

¹⁴GPs offers regular phone consultation hours (typically in the early morning).

¹⁵Emergency GP care was restructured in 2015 and thus there is a data break in the administrative data. Therefore, in our main analysis, we focus on GP contacts in the first four years of life where both treated and control children are exposed to the same regime of emergency GP care. Analyses that also include 2015 and later years (and only consider non-emergency GP care) lead to very similar results that are available on

independent, emergency GP contacts may be a more direct measure of poor health that requires attention. Importantly, we do not include child GP contacts in the preventive care program in our main outcome measure, but analyze those contacts separately. Thus our measures of GP contacts (scheduled contacts and emergency contacts) do not measure the participation in the voluntary preventive care program but focus on contacts due to health problems or parental concerns about the child's health. Moreover, our follow-up period of up to four years (and our analyses of GP contacts after the initial year of the child's life) allows us to speak to the role of substitution between nurse visits and GP contacts: While first-year effects on GP contacts may be caused by substitution, the scope for substitution in the longer-run is likely small.

As alternative measures of child health, we also consider two types of hospital contacts: Hospital admissions and outpatient contacts. Around 25 and 39 percent of children are admitted to the hospital or have an outpatient contact during their first year of life, respectively. While contacts to hospitals may capture more extreme health problems, these figures illustrate that, in general, hospital contacts are not rare and often related to routine check-ups. One aspect worth noting is that the 2008 strike covered all unionized nurses and thus hospital care for non-emergency patients was restricted. Therefore, GPs may have been more reluctant in referring children to hospitals in the strike period.

Second, we consider the impact of strike exposure on maternal postpartum mental health problems. These potential effects are interesting in their own right and also as mechanisms or reinforcing factors for longer-run effects of strike exposure on children. We create an indicator that is equal to one if mothers have any contact with a psychologist and/or psychiatrist in the first two years after the child's birth. We also consider the more extreme margin of maternal outpatient and inpatient contacts with psychiatric specialists up to two years after the child's birth.¹⁶

request.

¹⁶We include diagnoses (using the International Statistical Classification of Diseases and Related Health Problems (ICD) system) between F01-F99.

Third, we study the impact of strike exposure on parental health investment decisions. As we exploit information on a sample of children exposed to the nurse strike (and thus the absence of nurse visits at specific ages), we are constrained in our ability to use nurse registrations on parental inputs as outcome measures in our main analyses.¹⁷ Relying on administrative data instead, we consider indicators for participation in the GP preventive care program, participation in the vaccination program, and the timely completion of rounds in the vaccination program. As the nurse visits are closely spaced around the recommended age for the first year vaccinations, we assess whether missing a specific nurse visit impacts the probability of a timely vaccination, which we take as a proxy for parental health investments.

3 Empirical Methods

To examine the effects of the timing of NHV, we exploit children’s exposure to the nurse strike in a difference-in-differences framework. Specifically, we estimate the following reduced form relationship:

$$y_{it} = \alpha_0 + \sum_{j=-7}^{-1} \phi_j 1(\text{bin}30_{it} = j) \times 1(\text{Year}_t = 2008) \quad (1)$$

$$+ \sum_{j=-7}^{-1} \beta_j 1(\text{bin}30_{it} = j) + \gamma' \mathbf{X}_{it} + \boldsymbol{\lambda}_t + \epsilon_{it}$$

where y_{it} is an outcome measure, such as GP contacts in the first year of life for child i born at time t . In our analyses for outcome measures from the administrative data, we consider all children born in the 210 day period prior to April 15 in the years 2008, 2009 and 2010 (12,078 children).¹⁸ We split each period in seven 30-days bins and include indicators

¹⁷In supplementary analyses, we have constrained our sample to early strike-exposed children and study their outcomes at the nurse visit around eight months (D visit). We have considered indicators for nurse-observed issues concerning mother well-being, feeding, child-parent contact as well as indicators for any nurse comments at all and referrals by nurses. However, these analyses rely on a very small sample relative to the expected effect sizes (and the expected noise in the measurement of outcomes by nurses) and is thus not very informative. Unfortunately, the nurse data on infant feeding (duration of breastfeeding) in the archived data are of very poor quality and we cannot use them at all.

¹⁸As mothers given birth during the strike also had a larger probability of being discharged on the day

that are equal to one if child i 's date of birth is within a particular bin. We include a set of fixed effects for the relevant cohort, λ .¹⁹ The interactions of the period bins with an indicator for the 2008 cohort (the year of the strike) identify our estimates of interest: Children born prior to the strike in 2008 are treated while children born at the same dates in 2009 and 2010 are untreated. We omit the bin furthest from April 15 and children in this group constitute the reference group.

In our main specification, we include the following covariates (X_{it}): paternal and maternal total income, indicators for the highest level of education (primary school, higher education, university degree), indicators for currently studying and for being employed, an indicator for parental civil status (cohabiting, married) and indicators for missing covariates. All the X_{it} are measured one year prior to birth of the focal child. Additionally, we control for measures drawn from the birth records, including the number of prenatal midwife visits and indicators for parents being below 21 years old, indicators for having had a Caesarean section or a home birth, and indicators for the child having been low birth weight (below 2500g), a preterm birth (below 37 weeks), child gender and maternal smoking status at birth.

The coefficients from interacting the age bins and the strike period indicator provide intention-to-treat (ITT) estimates of strike exposure at a certain age relative to the reference group. To show that strike exposure is relevant, we present estimates for the impact of strike exposure on the probability of missing a nurse visit at a specific time in the child's life (the first stage). Furthermore, we present evidence on complier characteristics that substantiates our assessment of the strike as a broad treatment impacting families across many observable dimensions.

of birth and fewer midwife visits (Kronborg et al., 2016), including children born during the strike would confound the impact of NHV with the impact of other aspects of care.

¹⁹Note that the year indicators cross calendar years: As an example the indicator for the year 2008 (the treated year) is equal to one for all birth in the 210 days prior to April 15, 2008 and thus identifies births in the calendar years 2007/2008.

3.1 Identifying assumptions

For our estimates to identify the causal impact of exposure to the nurse strike, we make two identifying assumptions. First, we assume that, in the absence of the strike, the difference-in-differences between children born in specific periods up to April 15 in the strike and control years should be zero (common trend). Thus our framework allows for the years 2008, 2009 and 2010 to differ in levels. These differences could, for example, be due to overall trends in children’s health or macroeconomic shocks that affect care and health of children. Our focus on births from different months of the year also calls for a discussion of the impact of seasonality: We allow children born across seasons to be systematically different from each other (with respect to their average outcomes) as long as this seasonality is the same across all cohorts.

One way of empirically assessing the untestable common trend assumption is to study predetermined variables, which should be unrelated to treatment exposure. In other words, we estimate model (1) using parental and birth characteristics as dependent variables. Our treated and control groups are balanced across observable pre-treatment characteristics (Appendix Tables A2 and A3). We find very few differences across the groups and only at modest levels of significance.²⁰ Another informal test is the assessment of pre-trends in outcomes across groups. As we do not observe children’s GP visits prior to treatment, we consider maternal pre-birth outcomes: Appendix Figure A2 plots pre-birth averages of maternal GP contacts and mothers receiving medical contacts with a psychiatric diagnosis for the treated and control children (born in the 210 day period up to the strike in treated and control years).²¹ The figure shows similar trends and levels for both measures of maternal health prior to birth.

Second, we assume that there are no other policies or shocks that covary with the timing of the strike. To provide support for this assumption, we assess whether strike exposure

²⁰We have also tested the joint significance of the interaction between the age bins and the strike indicator in each of these regressions. None of the joint tests are significant at the 10 percent-level. Results are available on request.

²¹We include hospital contact diagnoses (using the ICD system) between F00-F99.

is related to differential health care provision through other channels than NHV. Similarly to Appendix Figure A2, Appendix Figure A3 plots the average number of prenatal midwife visits and GP consultations, the average number of days admitted to hospital after birth, and the share of mothers having a C-section for mothers in the strike-exposed year and control years. The graphs do not indicate systematic differences or trends in any of these types of care around birth across the groups that we consider.

A final concern that we address is individuals' selection out of the strike treatment or out of our sample. First, families could not to manipulate their treatment status since all children in our analysis sample were born either prior to the strike or a minimum of four months after the strike ended.²² Second, families could select out of our analysis sample by moving to a different municipality or out of the country. In our main analysis, to focus on children who were either treated with default care in Copenhagen or by the strike while residing in Copenhagen, we omit data for 1,962 children who move out of the municipality during their first year of life. If strike exposed families are more (or less) inclined to move, our estimates could be biased.²³ Appendix Figure A5 shows that this concern is not important as the share of children that we observe as Copenhagen residents during their first year of life is not impacted for treated and control cohorts. However, as a robustness check, we include domestic movers into our main analyses (so that only death and migration abroad cause exclusion).²⁴

²²In Appendix Figure A4 we show that the density of births around the strike does not indicate bunching around the beginning or end of the strike period.

²³As the strike was a nation-wide strike and of relative short duration (which parents were aware of), the risk of strike-induced domestic migration should be small.

²⁴We know individuals municipality of residence at January 1 each year. We restrict children born 210 days prior to April 15, 2008, 2009 and 2010 to still reside in Copenhagen at January 1, 2009, 2010, 2011 respectively.

4 Results

4.1 Descriptive Statistics

Table 2 presents summary statistics for our main sample of children born in Copenhagen across the groups of treated children (born September 18, 2007 - April 14, 2008) and children in the control group (born September 17, 2008 (2009) - April 14, 2009 (2010)). In the top panel, we present summary statistics for outcomes and covariates from the administrative data. In the bottom panel, we present variables on nurse visits from the nurse records from Copenhagen. In this panel, we further constrain our sample to the data periods in the years 2008 and 2009 as the nurse data is right-censored for the children born in 2010.

Control children have on average 4.6 and 10.4 GP contacts during the first and second year of life respectively. During the third and fourth year of life children have 10.2 contacts. Regular GP contacts constitute around two thirds of the total number of contacts. The infant vaccinations and preventive health checks have high coverage rates at around 90 percent. The treated and control groups are well-balanced across covariates.

Focusing on the bottom panel of Table 2, we find that the four universal nurse visits are well attended. The average number of universal visits per child is 3.3 for control children. This figure implies that the average child receives three out of the four universal visits. On average, children additionally receive one home visit scheduled due to a specific need. This average masks heterogeneity across children.

Table 2 Variable means, strike exposed and control period

	Treated group		Control group	
	Mean	Obs.	Mean	Obs.
Total GP 1st year	4.44	4081	4.55	8725
Total GP 2nd year	10.69	4049	10.35	8649
Total GP 3-4 years	11.10	3955	10.22	8451
Emerg. GP 1st year	1.42	4081	1.47	8725
Emerg. GP 2nd year	3.75	4049	3.46	8649
Emerg. GP 3-4 years	3.50	3955	3.18	8451
Vacc., 1st round	0.85	4081	0.90	8725
Vacc., 2nd round	0.87	4081	0.91	8725
Vacc., 3rd round	0.88	4081	0.91	8725
Prev. care, 5 weeks	0.88	4081	0.92	8725
Prev. care, 5 months	0.92	4081	0.93	8725
Prev. care, 12 months	0.93	4081	0.93	8725
Emerg. GP 1st year mothers	0.72	4081	0.70	8725
Emerg. GP 2-4 years mothers	2.10	3950	1.97	8445
Mother psych. diag. 1st year	0.01	4081	0.01	8725
Mother psych. hosp. adm. 3 years	0.01	4081	0.01	8725
Mother psych. outpat. cont. 3 years	0.03	4081	0.03	8725
Midwife visits	4.80	3970	4.75	8507
Smoking status, Mother	0.10	4014	0.09	8587
Child sex	0.48	4081	0.48	8725
Low birth weight	0.04	4009	0.06	8598
Preterm birth	0.06	4014	0.06	8587
C-section	0.21	4081	0.21	8725
Home birth	0.01	4081	0.01	8725
Cohabiting	0.76	4081	0.78	8725
Married	0.37	4081	0.39	8725
Prim. school, mother	0.15	4081	0.12	8725
Uni. degree, mother	0.30	4081	0.32	8725
Student, mother	0.05	4081	0.05	8725
Employed, mother	0.77	4081	0.77	8725
Danish, mother	0.76	4081	0.74	8725
Young mother	0.02	4081	0.02	8725
Young father	0.01	4014	0.01	8551
Income, mother	281.78	4081	289.58	8725
No. of nurse visits	3.77	4081	4.40	4269
Number of registered A-D visits	2.70	4081	3.28	4269
No initial visit	0.16	4081	0.08	4269
No 2-month visit	0.44	4081	0.25	4269
No 4-month visit	0.44	4081	0.24	4269
No 8-month visit	0.26	4081	0.15	4269

Notes: The sample includes children who were born in Copenhagen in the treated period (September 18, 2007 - April 15, 2008) and in control periods (September 17, 2008 and 2009 - April 15, 2009 and 2010). For the data from the nurse records (bottom panel), the control group only includes the period September 17, 2008 - April 15, 2009.

Table 2 also illustrates the impact of strike exposure on the program coverage: For all types of visits, treated children have a higher probability of missing the given visit. The difference in the number of universal visits across groups is identical to the difference in their total number of visits. This finding indicates that the average number of extra visits was not affected dramatically by the strike. In the following, we will analyze these patterns in greater details.²⁵

4.2 First Stage and Compliers

In Denmark, both private and public wages are to a large degree determined by collective bargaining (Ibsen et al., 2011). In 2008, the negotiations for all publicly-employed nurses, midwives and a large fraction of other employees in the public health sector broke down and resulted in a conflict. Thus on April 15, 2008 the unionized employees in the health care sector went on a national strike. As a result, a total of 45 percent of public employees were on strike in the following weeks (Due and Madsen, 2008). The strike lasted 61 days and ended on June 14, 2008.²⁶ During the strike period, only managing nurses and a small fraction of regular nurses (employed on specific terms and thus not participating in the strike) were on duty in Copenhagen, the setting for our analysis. These nurses carried out around one tenth of the expectable non-strike default of nurse visits.²⁷ Moreover, they provided phone services for families that were affected by the strike.

²⁵To assess the representativeness of our sample of families from the capital of Denmark, Appendix Table A4 compares children and parents from Copenhagen to the general Danish population of parents. Children and parents from Copenhagen differ from the general population on a number of characteristics: they are more likely to cohabit and less likely to be married. Mothers from Copenhagen have a higher educational attainment. Parents from Copenhagen are less likely to be employed and of Danish origin. With respect to children's health and characteristics, children in Copenhagen resemble children from the rest of county: 5 percent of children are low birth weight children and 7 percent are born prematurely. Children in Copenhagen are also similar to the rest of Denmark with respect to the number of nights at hospital after birth, the number of prenatal midwife visits, the rate of C-section deliveries, and the share of home births. At the same time, 62 percent of children born in Copenhagen are firstborns compared to 43 percent outside Copenhagen, their parents are older and less likely to smoke.

²⁶The unions demanded a 15 percent wage growth. The agreement resulted in a 13.3 percent wage increase over a three-year duration.

²⁷We calculate this share of performed visits by comparing the strike period to the same period in the following year.

Appendix Figure A1 presents graphically the impact of strike exposure on the number of nurse visits for children in the treated and control cohorts in Copenhagen (2007/2008 and 2008/2009, respectively). Strike exposure impacted the total number of nurse visits that children received. Panel (a) of Appendix Figure A1 shows that control children receive an average of 3.3 visits while treated children receive 2.7. Panel (b) shows the total number of visits (universal + extra) divided by treatment status. The youngest strike exposed children appear to not only lose one but two nurse visits. This finding reflects that early hospital-discharged children receive two visits within the first 14 days of life - one universal visit and one extra visit. In section 4.5, we examine the robustness of our general conclusions to the omission of this group of children (a doughnut hole-approach).

To further examine the impact of the strike on nurse visits and to illustrate the identifying variation that we use (i.e., the overall decrease of the number of visits is driven by a lack of visits at specific ages), Figure 2 shows the impact of strike exposure on the probability of missing a specific nurse visit. The figure shows the raw relationship between date of birth and missing a nurse visit estimated with kernel weighted local polynomials. We use an epanechnikov kernel, a rule-of-thumb bandwidth and 42 (5-day) smoothing points throughout. Black lines and confidence intervals are for the treated period, grey lines and confidence intervals are for the control periods.²⁸ The graphs plot the probability of missing a nurse visit for children born in the 210 days before the strike for the years 2008 and 2009.

In absence of strike, the share of children, who miss a specific nurse visit, is stable as indicated by the grey lines in Figure 2. 60 percent of children born immediately before the strike miss the initial visit while all children older than approximately 20 days at strike start miss the initial visit with unaffected probability (20 percent). Panels (b) and (c) show that missing the two and four month visits is also correlated with child age at strike. Finally, only the oldest children in our sample have an increased probability of missing the eight month visit while all the younger children are unaffected at that time (because the strike ended by

²⁸We construct graphs that plot outcomes similarly unless otherwise noted.

the time their visit was due).




Fig. 2 Share of children with missed nurse visits for children born in the treated and control period
Notes: The figure shows the raw relationship between date of birth and missing a nurse visit estimated with kernel weighted local polynomials using an epanechnikov kernel, a rule-of-thumb bandwidth and 42 (5-day) smoothing points. The black line and dashed black confidence intervals are for the treated period, the grey line and dashed grey confidence intervals are for the control period. Treated period: September 18, 2007 - April 15, 2008. Control period: September 17, 2008 - April 15, 2009).

Table 3 presents formal estimates from regressions based on Equation (1). Coefficients reflect the effect of being born in a specific bin on the probability of not receiving each nurse visit (the omitted baseline is the 30 days bin furthest from strike start). The columns show results for the different types of universal nurse visits. The regression results mirror the graphical representation: The strike only has an impact on the initial visit for children who

were between 30-0 days at strike start. On average children in this bin have 17.1 percent-points higher probability of missing the initial visit (relative to the reference group). Children who were 90 days and below at strike start have an increased probability of a missed two month visit with the 60-31 bin most severely affected (51.1 percent-points). Children who were between 61 and 150 days at strike start have their four month visit most severely affected by the strike. Only the oldest children in the strike exposed period have increased probability of a missed eight month visit compared to younger children (around 40 percent-points difference when compared to the children, who were youngest at strike start). As shown in column (5) strike exposure does not differentially impact children’s number of completed universal visits. However, children in the 30-1 day bin loose on average 0.267 nurse visits more than the reference group (significant at the 10 percent level). This result reflects that children below age two weeks at strike start potentially loose two visits, the universal initial visit and an additional early visit if discharged shortly after birth.

Having established that age at strike start has a meaningful impact on timing of the missed nurse visit for strike-exposed children, we have the concern that nurses strategically chose the children they visited, i.e. that only the most well-off children were impacted by the child. This question is important for the interpretation of our findings. In general, the large scale of the strike—with only one tenth of performed nurse visits in Copenhagen during the strike relative to the default—suggests that the strike impacted large parts of the population. However, our unique data also allows us to characterize compliers (i.e. children who missed nurse visits due to the strike) more formally in our sample.

Table 4 characterizes the compliers with respect to the probability of missing the first nurse visit (analyses for the other three universal visits lead to similar conclusions and are available on request). Following Angrist and Pischke (2008), we characterize the compliers by i) splitting the full sample into relevant subgroups, ii) estimating the model for each subgroup individually and iii) calculating the ratio between the coefficients from each subgroup and the full population. The ratios are the relative likelihood that a complier belongs to that

Table 3 First stage: Effects of strike exposure on the probability of a missed nurse visit

	(1)	(2)	(3)	(4)	(5)	(6)
	No initial visit	No 2-months visit	No 4-months visit	No 8-months visit	Number of registered A-D visits	No. of nurse visits
Days						
180-151	0.002 (0.026)	-0.040 (0.037)	0.100*** (0.037)	-0.324*** (0.034)	0.261*** (0.091)	0.223 (0.166)
150-121	0.003 (0.026)	-0.018 (0.037)	0.247*** (0.037)	-0.357*** (0.034)	0.126 (0.090)	0.205 (0.162)
120-91	-0.027 (0.026)	-0.017 (0.037)	0.364*** (0.037)	-0.363*** (0.033)	0.043 (0.088)	0.181 (0.164)
90-61	-0.007 (0.025)	0.155*** (0.038)	0.225*** (0.038)	-0.353*** (0.034)	-0.020 (0.087)	0.247 (0.163)
60-31	-0.005 (0.024)	0.511*** (0.035)	-0.039 (0.036)	-0.423*** (0.033)	-0.044 (0.083)	0.115 (0.153)
30-1	0.171*** (0.028)	0.323*** (0.037)	-0.079** (0.036)	-0.395*** (0.033)	-0.019 (0.085)	-0.267* (0.158)
Obs.	7874	7874	7874	7874	7874	7874

Notes: Each column shows estimates from separate regressions. The coefficients are for the interactions of 30-day bins and a strike year indicator. All regressions include period and bin fixed effects, as well as control variables. Parental covariates are paternal and maternal income, indicators for the highest level of parental education (primary school, high school, university degree), indicators for the mother currently studying or being employed, parental cohabitation and marital status and separate indicators for missing covariates. All covariates are measured one year prior to birth of the focal child. Child/birth covariates include indicators for parental age below 21 at birth, indicators for a C-section, home birth, low birth weight (below 2500g), a preterm birth (below 37 weeks), child gender, maternal smoking status at birth and the number of prenatal midwife visits. The sample includes children born in Copenhagen in the treated period (September 18, 2007 - April 15, 2008) and in control period (September 17, 2008 - April 15, 2009). The outcomes in columns (1)-(4) are indicators for the probability of having missed the respective universal home visit. The outcome in column (5) is the number of universal nurse visits received. Column (6) presents results for the total number of nurse visits (universal and additional visits). Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

particular subgroup. We look at the first stage estimates across groups of families defined by characteristics that may at least be partly observed by the nurses: child gender, parental education in a health-related field,²⁹ initial child health,³⁰ and child parity. We show coefficients for the 30-day bin as only children born in that bin were affected by the strike. In general, the complier analysis suggests that the strike affected the considered subgroups relatively similarly and a stronger first stage does not covary with characteristics that may indicate positive potential outcomes. Thus we think it is reasonable to state that nurses did not prioritize to a great degree based on the given characteristics. This finding is relevant for our interpretation of especially heterogeneous effects (because we can rule out that nurses' prioritized certain subgroups during the strike as a main driving factor).

Taken together, strike-exposed children missed on average one nurse visit. Thus we cannot fully disentangle the effect of having one less nurse visit from the effects of timing. Strike exposed children missed this visit at different ages and we compare outcomes of children across years relative to the reference group of children born 180-210 days prior to the strike. Our first stage results provide powerful evidence for the differential timing of the assigned treatment (one less visit). Thus we think it is reasonable to interpret our findings as predominantly being driven by timing given that the different visits coverage-specific topics, as outlined in section 2.

4.3 Main Results: Child and Maternal Health

To measure the impact of strike exposure at different ages on children's and mother's health, we use outcomes from the administrative data. Figure 3 presents graphical evidence of the raw relationship between age at strike start and accumulated GP contacts at ages one through four.³¹ The number of accumulated GP contacts reveal a clear pattern: Children,

²⁹Having parents with an educational background in a (child) health-related field implies that either one of the parents are educated as doctor, midwife, nurse or pedagogue.

³⁰We define a children with low initial health as having a birth weight below 2500g and/or being born preterm.

³¹Figures for regular and emergency contacts are available on request.

Table 4 Compliers: Effects of strike exposure on the probability of missing the initial visit by subgroup

	Gender		Education		Initial health		Parity	
	Boys (1)	Girls (2)	Not health (3)	Health (4)	Not poor (5)	Poor (6)	>1 (7)	=1 (8)
30-1	0.199*** (0.040)	0.130*** (0.039)	0.151*** (0.032)	0.217*** (0.059)	0.157*** (0.029)	0.237** (0.117)	0.132*** (0.046)	0.194*** (0.035)
Ratio to full pop.	1.19	0.78	0.90	1.29	0.94	1.41	0.78	1.15
Control group mean	0.09	0.07	0.08	0.09	0.08	0.11	0.10	0.07
Observations	4101	3773	6156	1718	7276	598	3026	4848

Notes: See notes for Table 3. In this table, we present estimates for the interactions of 30-day bins and a strike indicator from separate regressions for various subgroups along with the ratio between the full-sample estimates and the various subgroup-estimates (both sets of regressions excluding control variables). We only show the estimates for the 30-1 day bin because only children in this bin had their initial visit affected by the strike in the full population. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

who were youngest at strike start in 2008 have significantly more GP contacts relative to children of older age groups and this pattern looks different in the control group. As Figure 3 further illustrates, there is a gradient inside the early strike-exposed group of children such that the youngest children have most GP contacts. This finding indicates that earlier NHV is relatively more important for child health than later NHV. For children older than 100 days at strike start, the average number of GP contacts is similar to the average for control children. Interestingly, the impact of missing an early nurse visit is persistent as the differences increase as the children ages.




Fig. 3 Accumulated number of GP contacts for children born in the treated (September 18, 2007 - April 15, 2008) and control periods (September 17, 2008 and 2009 - April 15, 2009 and 2010)
Notes: The figure shows the relationship between date of birth and accumulated total GP contacts. See Figure 2 for further details.

Table 5 shows our main results for the impact of strike exposure on child health decomposed by the type of GP contact. To rule out that substitution toward GP visits during the first year of life drive our findings, we present estimates for *yearly* outcome measures, i.e., child GP contacts measured in each year of life of the child.³² Across periods the estimated effects are significant and the patterns documented in the graphical analyses persist: Children born in the 30-1 days age group have 1.8, 1.6 and 1.3 additional GP visits during the first, second and third to fourth year of life. In percentage terms (evaluated at the average number of GP visits of the control group) our results translate to 40.3 percent, 15.5 percent and 12.4 percent increases. Considering emergency GP contacts, the relative effects are larger at 50.0 percent, 18.4 percent and 18.2 percent during the first, second and third to fourth year of life. Children in the 60-31 days age group have significantly more GP contacts (across types) in their second year of life. For all other age groups the timing of strike exposure has no significant effects on GP contacts.

To assess the impact of strike exposure at other margins, we have also considered alternative measures of child health (Appendix Table A5): child hospitalizations and outpatient contacts. While most point estimates for first year hospitalizations are imprecise, we find suggestive evidence that early strike-exposed children are 7-8 percent-points (40 percent) more likely to be hospitalized during the second year of life. These results carefully support that our results for GP care and indicate actual health effects that do not exclusively reflect substitution and precautionary parental behavior. Furthermore, we see some indication for a decrease in first year outpatient contacts. While nurses in non-strike years can refer families as outpatients to hospitals in case of health or feeding issues, during the strike this option

³²We have also estimated the regression equivalents of the graphs for the *accumulated* GP contacts for all years between year one and four in one combined graph. The effects on GP contacts increase as the child ages, in particular during the first two years of life. At age four, treated children have 4.6 (18.3 percent) more GP contacts in total for the 30-1 bin, 2.8 (11.1 percent) for the 60-31 bin and 2.4 (9.5 percent) for the 90-61 bin. For regular GP contacts the percentage effect is 15.8 percent for the youngest age groups and 8.1 percent for the 60-31 age bin (for the 90-61 age bin we see no significant effect on the number of regular GP contacts). The percentage effects on emergency contacts are 23.2 percent, 17.1 percent and 13.4 percent for the 30-1, 60-31 and 90-61 age bins.

Table 5 Effects of strike exposure on child health: Yearly GP contacts by type

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
	Total GP 1st year	Total GP 2nd year	Total GP 3-4 years	Ordin. GP 1st year	Ordin. GP 2nd year	Ordin. GP 3-4 years	Emerg. GP 1st year	Emerg. GP 2nd year	Emerg. GP 3-4 years
Days									
180-151	0.469 (0.516)	0.129 (0.515)	0.707 (0.597)	0.142 (0.348)	-0.147 (0.319)	0.447 (0.391)	0.327 (0.222)	0.277 (0.296)	0.260 (0.312)
150-121	0.297 (0.517)	0.402 (0.508)	0.539 (0.594)	0.149 (0.342)	0.191 (0.319)	0.358 (0.386)	0.148 (0.229)	0.211 (0.290)	0.181 (0.310)
120-91	-0.187 (0.527)	-0.161 (0.492)	0.007 (0.564)	-0.212 (0.355)	0.022 (0.310)	0.193 (0.383)	0.025 (0.222)	-0.183 (0.283)	-0.186 (0.291)
90-61	0.758 (0.550)	0.701 (0.512)	0.849 (0.580)	0.332 (0.379)	0.445 (0.329)	0.509 (0.390)	0.426* (0.223)	0.255 (0.284)	0.340 (0.303)
60-31	0.364 (0.529)	1.707*** (0.508)	0.692 (0.576)	0.087 (0.352)	0.923*** (0.321)	0.408 (0.381)	0.277 (0.227)	0.783*** (0.291)	0.284 (0.300)
30-1	1.835*** (0.555)	1.614*** (0.519)	1.271** (0.584)	1.109*** (0.376)	0.977*** (0.332)	0.692* (0.394)	0.726*** (0.233)	0.637** (0.288)	0.579* (0.296)
Control group mean Obs.	4.55	10.35	10.22	3.09	6.89	7.04	1.47	3.46	3.18
	12078	11982	11709	12078	11982	11709	12078	11982	11709

Notes: See notes for Table 3. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

was likely limited (due to nurses in hospitals also being on strike).³³ Given that we do not see longer-run impacts of strike exposure on outpatient contacts, we conclude that our finding for outpatient care supports the idea of some substitution of care during the strike (from hospital care to GP care).³⁴

Our main results show that early strike-exposure impacts children’s number of GP contacts – in the short and longer run. Importantly, nurses also focus their attention on maternal physical and mental well-being. Table 6 presents results for maternal (total and emergency) GP contacts, as well as (non-emergency) maternal contacts to psychologists and psychiatrists (after referrals from their GP). Finally, while we also consider maternal psychiatric hospital admission and outpatient contacts within the first two years after birth. However, this outcome is a very rare event limiting our ability to detect impacts given our design and sample size.

Table 6 shows that mothers, who are strike-exposed shortly after the birth (90-1 days) of their child, have 1.8-2.6 additional GP contacts (9.5-13.6 percent increase at the mean) during the second to fourth year of life but no additional visits in the first year. Also for mothers, the GP results are both driven by scheduled and emergency GP contacts. For our measure of contacts to a psychologist or psychiatrist two years after birth, we find that mothers with early strike exposure (30-1 days bin) are 3.6 percent-points more likely to have a contact with a specialist (72 percent). In sum, our results indicate that early strike-exposure that resulted in reduced access to early NHV has impacts also on maternal psychical and mental

³³However, hospitals were obliged to ensure an adequate level of care provision.

³⁴We have also attempted to analyze child outcomes based on nurse registrations at age eight months and longer run outcomes: Constraining our sample to children who received the eight month visit, we do not find precise estimates for the impacts of strike exposure on child development at eight months. However, these analyses are based on around 40 percent of our main analysis. Considering longer-run outcomes, we have explored the impact of timing of strike exposure on the probability of delayed school start of children. We do not detect any effects. Given the age of the strike-exposed and control children, we cannot yet examine longer-run impacts of the 2008 strike on academic test scores (observed for the first time during grade two). Assessing the school entry examination of around 75 percent of the children in our sample, we do not see any impact of timing of strike exposure on child BMI or probability of being overweight. In our sample we likely lack power to analyze these outcome (given low level of obesity prevalence at around 7 percent). Furthermore, we miss 25 percent of children in our school entry records that only cover Copenhagen and thus do not include children, who move.

Table 6 Effects of strike exposure on maternal health: GP contacts, postnatal psychiatric diagnoses and contacts with psychiatric specialists

Days	(1) Total GP 1st year mothers	(2) Total GP 2-4 years mothers	(3) Emerg. GP 1st year mothers	(4) Emerg. GP 2-4 years mothers	(5) Psychologist psychiatrist two years	(6) Psych. hosp. adm. and output. cont. two years
180-151	-0.087 (0.375)	-1.325 (0.986)	0.007 (0.097)	0.046 (0.217)	0.025 (0.016)	-0.002 (0.002)
150-121	0.024 (0.371)	1.205 (0.984)	0.067 (0.103)	0.204 (0.210)	0.024 (0.015)	0.003 (0.003)
120-91	0.439 (0.394)	0.802 (1.013)	0.015 (0.098)	0.250 (0.215)	-0.001 (0.015)	-0.000 (0.002)
90-61	0.366 (0.390)	2.355** (1.019)	0.124 (0.096)	0.633*** (0.225)	0.025 (0.016)	0.005 (0.003)
60-31	0.274 (0.399)	1.862* (1.034)	0.215* (0.121)	0.702*** (0.228)	0.019 (0.016)	0.008** (0.003)
30-1	0.506 (0.391)	2.664*** (1.007)	0.124 (0.097)	0.585*** (0.217)	0.036** (0.016)	0.001 (0.002)
Control group mean	7.29	19.92	0.70	1.97	0.05	0.00
Obs.	12078	11698	12078	11698	11982	11982

Notes: See notes for Table 3. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

health. Moreover, effects on maternal well-being may constitute a mechanism for or reinforce the health effects on children that we have documented.

4.4 Mechanisms

Our main analyses show that early strike-exposure matters for child and maternal health. We interpret this finding as support for the hypothesis that early NHV matters more for the considered health outcomes than later visits. To speak to potential mechanisms for the observed effects, we focus on the elements of the composite nurse treatment that are of particular importance in the initial visits: information and counseling, and screening and monitoring of infant and maternal health. First, to assess the importance of information and counseling in explaining the negative effects of forgoing an early nurse visit, we study heterogeneous effects across two relevant dimensions: the parity of the child and parental health-related education.³⁵ Specifically, we hypothesize that first-time parents and parents without professional knowledge about child health and development may see larger effects of early strike exposure if information is an important element that strike-exposed parents lack.

For brevity, we present results for our measure of total GP contacts in year one and year two through four of the child’s life.³⁶ We split our sample into subgroups and additionally estimate a fully interacted model on the full sample. Appendix Table A6 presents our split-sample results.³⁷ Column (1)-(4) show regression results for samples divided into groups of parents with and without an education in a health field. While we do not find significant effects of the timing of strike-exposure for children of parents educated in a health-related field, for children of parents *not* educated in those fields, our results resemble the main results. Similarly, first-born children see stronger effects of early strike exposure and a larger gradient than higher parity children as shown in column (5)-(8) in Appendix Table A6. While

³⁵The group with parents educated in health include children who have at least one parent educated as either a medical doctor, midwife, nurse or pedagogue.

³⁶Results for emergency GP contacts only are available on request.

³⁷Appendix Figure A6 presents the raw relationship between the timing of strike exposure and GP contacts accumulated at age four divided by parental health education and parity.

we formally cannot reject the null hypothesis that the effects are the same across subgroups (see Appendix Table A7), our findings suggest that an information and counseling channel is important for explaining longer-run health impacts of early NHV. At the same time, as illustrated in Appendix Table A8, we find less systematic differences in estimates across families of high or low socio-economic status, if anything, high SES families appear to see larger effects of early strike exposure.³⁸ This finding may further underline the importance of specific guidance and information for new parents and, additionally, points to the potential importance of another channel for early life NHV, namely universal screening and health monitoring.

Early NHV puts a focus on screening for potential health problems in infants and mothers: Offered as a universal program, it represents an early window of opportunity to detect and confront health problems. Our results for maternal mental health in Table 6 suggest that lack of early screening negatively impacts maternal mental health. Another way of further examining the importance of screening is to assess the performance of nurses with respect to screening in non-strike years.

Figure 4 presents nurse registrations, referrals and maternal health care usage for two groups of mothers in our control year data: first, mothers with registrations of maternal mental health problems at the initial visit (10 percent of mothers) and without those registrations. Conditional on having follow-up visits, we observe interesting patterns that point to the importance of nurse screenings very shortly after birth: Nurses are more likely to register mental health problems in later visits for early-detected mothers. Additionally, mothers with early detected mental health problems receive more referrals to other health professionals and, importantly, among early-detected mothers there is a higher prevalence of externally measured mental health issues.

³⁸Appendix Table A8 also examines heterogeneity by gender, child initial health, and parental risky behaviors (proxied by maternal smoking during pregnancy). We see indication for boys, children with poor initial health and children of parents with risky parental behavior being relatively more affected by the absence of early NHV (however, also in these analyses, we cannot reject equality of effects in most cases).




Fig. 4 Share of mothers with nurse registrations, referrals and contacts to psychologists/psychiatrists by registered maternal well-being concerns (0/1) at initial nurse visit, mothers of children born in the control period.

Notes: This figure divides mothers of control children (born in Copenhagen between September 17, 2008 - April 15, 2009) into two groups: The 10 percent of mothers with nurse registrations in their initial nurse visit (concerned about maternal well-being vs not concerned) and the 90 percent of mothers without a registered concern. We constrain the control sample to mothers, who received the initial visit and plot the share of mothers who receive registrations of maternal well-being issues at later visits (B, C and D), the share who are referred to other health care professionals by nurses, and the share for whom we observe any contacts with psychologists/psychiatrists up to two years after their birth.

Relating Figure 4 to the overall prevalence of maternal mental-health related contacts, our calculations suggest that nurses during their first visit identify up to one out of four of those mothers who end up having a mental-health related contact with specialists in the first two years of their child’s life.³⁹ This illustrative figure suggests large potential health returns from early screening efforts.

A final and important potential pathway for the effect of early NHV are parental investments in response to those. Nurses provide information and guidance about issues such as

³⁹Nurses screen around 10 percent of mothers in the sample as having a mental health problem. Of those, 13 percent end up having at least one psychologist/psychiatrist contact in the first three years of the child’s life. In the population, the prevalence of those contacts is around 5 percent. These figures suggest that nurses may capture around 20 percent of those mothers, who end up with a contact.

other available health care services, appropriate interactions with children at different ages, and aspects such as sleep and child feeding. However, given our sample size in combination with our empirical strategy, we are constrained in an analysis of those parental behaviors: Appendix Tables A9 and A10 study whether strike exposure impacts participation in the childhood preventive care program and vaccination program participation as outcomes.⁴⁰ As the tables illustrate, we cannot draw firm conclusions due to very imprecise estimates.

4.5 Robustness Tests

Our main results are robust to a number of changes to our main specification and sample. For brevity, we only present robustness tests using our yearly measures of child GP contacts as outcome in the appendix material. We show that our conclusions are not sensitive to the omission of individual-level control variables (Appendix Table A11) and reasonable alternative choices of bin size (Appendix Tables A12 and A13). To rule out that our measure of strike exposure captures other factors, we implement a set of placebo regressions: Appendix Table A14 shows estimates from those regressions where we define “treated” children as those born 210 days prior to April 15, 2009 (the year after the strike). We find no significant effects of strike exposure in the placebo regressions.

In additional robustness tests, we have ruled out that including movers from Copenhagen alters our conclusions and confirmed that they are robust to the implementation of a doughnut hole approach (where we drop children born within 20 days of strike start, who were likely to lose more than one visit on average). By using earlier cohorts of children as a control group, by examining the impact of strike exposure on children aged five during the strike, and by constraining our main analysis to using data from the years 2008 and 2009 (our “first stage” sample), we confirm that our choices of control and treatment groups do

⁴⁰ Almost 80 percent of children receive all infant vaccinations and each round of vaccinations are attended by 90 percent of children in Copenhagen. Participation in the vaccination program is voluntary and the decision ultimately rests at the parents. The DNBH specifically mentions nurse visits as a central strategic element to promote the benefits of vaccinations to parents (The Danish National Board of Health [Sundhedsstyrelsen], 2018). The DNBH report highlights the close relationship between the families and their assigned nurse which facilitates dialogue if parents are in doubt or have chosen not to participate.

not drive our findings. Our main conclusions—that earlier strike exposure is relative more important for children’s health—remain intact across these iterations.⁴¹

5 Costs and Benefits

In this section, we perform a stylized analysis of immediate health benefits and the costs of early NHV (relative to later NHV). Specifically, we relate the value of prevented GP visits for mothers and children to the costs of those visits. Given the most consistent evidence for an impact of the strike on the health of children and mothers exposed early, we focus in the following on the initial and two-month nurse visits. The assessment of the benefits of early visits is—due to our design—always relative to the benefits of later visits. Put differently, in our calculations, we assume that the the benefits of the later visits are zero.

Benefits Appendix Table A15 presents results for the impact of strike exposure on GP fees (for both mother and child) at age four.⁴² As we disregard longer-run benefits, such as prevented child hospital admissions, and potential spill-over effects to other domains, such as child cognitive development or maternal timely return to the labor market, our measure of benefits (prevented GP costs) is likely very conservative.

Children born in the 30-1 and 60-31 days age groups (and their mothers) have significantly higher GP expenses, in line with our finding of increased GP contacts for these groups. Specifically, children and mothers impacted by the strike in the given groups have, respectively, 154.3 and 94.2 EUR higher GP expenses at age four. To translate these costs (or the benefit from preventing them) into a measure directly linked to a forgone visit, we scale the reduced form estimates with the probability of missing the specific visits for the given groups of children and mothers.⁴³ Thus we estimate the benefits of the initial nurse visit and the two

⁴¹All mentioned robustness tests for yearly GP and other outcomes are available on request.

⁴²GPs are reimbursed for all procedures they provide to patients in a given calendar week. We do not find clear evidence for the treated children having more costly GP visits on average.

⁴³For the first group (children born 30-1 days prior to strike) both the probability of not receiving the initial and two-months visits are increased by 17.1 and 32.3 percentage points, respectively (see Table 3). Thus

months nurse visit as 554.2 EUR and 184.3 EUR (prevented GP costs for child and mother up to the child’s four year birthday).

Costs To quantify costs of a home visit, we only consider the direct costs related to nurses’ salaries.⁴⁵ Additionally, we assume that all types of home visits have the same average cost. We calculate the cost of a home visit in two different ways that allow us to bound our calculations: first, we conservatively assume that municipal nurses spend all working time on home visits. Second, in the alternative scenario we incorporate that nurses have other tasks beyond home visits (such as supervision of school children, consultancy and phone hours, team meetings, administrative tasks).

We estimate the weekly number of canceled visits during the strike to be 760.⁴⁶ After the strike, the municipality of Copenhagen reported daily savings during the strike of 35,500 EUR per workday or 177,500 EUR per (business) week (because the municipality did not pay salaries to the unionized nurses on strike). For our most conservative measure of costs per visit, we divide the weekly divided by the weekly number of canceled visits, $177,500 \text{ EUR}/760 \text{ visits} = 233.6 \text{ EUR per visit}$. For our alternative measure—that takes into account that nurses also have other obligations—we adjust the share of working hours nurses dedicate to home visiting to 55 percent.⁴⁷ Dividing the weekly savings during the strike adjusted with the actual time spent on home visits by the number of canceled visits,

to calculate the benefit of the initial visit, we scale the increase in GP fees for the 30-1 day group with the increase in their risk of missing the initial visit while subtracting the share of their increase in GP fees that can be attributed to the higher probability of also missing the two month visit: $(154.3 - 184.3 \times 0.323)/0.171 = 554.2 \text{ EUR}$.⁴⁴ For the 60-31 day age group only the probability of missing the second nurse visit was impacted by the strike (51.1 percentage points). Thus, we scale their increase in GP fees due to strike exposure with the increase in the risk of forgoing the two month visit: $94.2/0.511 = 184.3 \text{ EUR}$.

⁴⁵We abstract from any fixed and variable costs beyond salaries to nurses. Examples of fixed costs are the education of nurses, capital (cars, building stock and software). Variable costs beyond salaries to nurses are management costs, cleaning services, transportation, lunch and coffee among others.

⁴⁶In our nurse data we observe that, during the full seven weeks of the strike, 85 weekly nurse visits were preformed. In the equivalent weeks of the following year, the weekly average of visits was 845. We assume that the difference in weekly visits equals the number of canceled visits caused by the strike ($845 - 85 = 760$).

⁴⁷In our data for the control period, 155 nurses performed visits implying that the average nurse had $845/155 = 5.5$ weekly visits. Assuming that one visit lasts 1.5 hours and that nurses spend an additional 1.5 hours on preparation, transportation and registration, nurses spend $5.5 \text{ visits} \times 1.5 \text{ hours at actual visit} \times 1.5 \text{ hours for tasks related to visit} = 16.5 \text{ hours weekly on NHV}$. If we assume that the average nurse work 30 hours per week, we estimate that nurses spend $16.5/30 = 55$ percent of their working time on NHV.

we find that the cost of a home visit in our alternative scenario is 128 EUR.⁴⁸

Comparing costs and benefits In both described scenarios for our calculations of costs, the initial nurse visit has a positive return of between 330.6 and 425.7 EUR. This represents a substantial return given that we only included savings related to GP care and under the fairly conservative assumption that the four month and eight month visits have zero benefits. For the two month visit, we conclude that the return only related to prevented GP costs is between -39.3 and 55.8 EUR. Thus our simple analysis indicates that early universal NHV is a cost-effective intervention. Our estimates highlight the importance of timing: While the cost of an initial visit is considerably lower than the associated health care savings at age four, the difference in the increase in GP fees and the savings from canceling a two month visit is considerably smaller.

6 Conclusion

Using nurse records linked to administrative data and exploiting exogenous variation induced by a national strike, we provide causal evidence on the impact of timing of early life health investments on child and maternal health. Thus our analysis of the effects of NHV moves beyond the extensive margin of treatment exposure. Studying the Danish universal program, we find that early NHV (during the initial weeks and first two months of the child’s life) impacts both child and maternal health trajectories (measured in our analyses as health care usage). We conclude that earlier visits are more important for children’s and mother’s health than later visits. While we cannot fully disentangle underlying reasons for increased health care usage for children and their mothers, we show that access to early NHV impacts emergency GP contacts and children’s hospitalization—also when we omit first year outcomes. Both findings point to actual health effects rather than substitution.

The heterogeneity of effects by parental health knowledge and child parity point to the

⁴⁸ $(177.500 \text{ EUR} \times 55\text{percent})/760 \text{ visits} = 128 \text{ EUR per visit}$

importance of information and parental confidence as a channel for health effects—supporting both is at the core of early home visits. While we do not directly observe parental beliefs and only have few measures of actual parental investment behaviors, both factors may be contributing to the effects of early home visits that we find.

Importantly, indicating the importance of timely screening for child and maternal health issues, we find that early NHV also plays a role for maternal postpartum mental health. As a consequence, our results imply that early home visits are likely to impact children through their impact on mothers: Existing research documents strong correlations between maternal postnatal mental health and child outcomes in different domains, and highlights the importance of early detection of maternal mental health problems. Thus early universal home visits can play an important role in securing population maternal and child health through the prevention of undetected and hence untreated mental health problems. In this aspect, our study echoes the finding of other recent work pointing to the importance of supporting the health of new mothers.

Finally, while initial visits in the Danish program focus on mother and infant physical health, infant feeding and sleep patterns, and maternal mental well-being, later nurse visits increasingly focus on other and more diverse domains of child development and parent-child interactions. In our setting, we do not find that those later visits impact the health outcomes that we can study. However, these visits may play an important role in further shaping parental investments and child development in other domains. We leave this topic as an important alley for future research.

References

- Almond, D. and J. Currie (2011). Killing me softly: The fetal origins hypothesis. *Journal of Economic Perspectives* 25(3), 153–72.
- Almond, D., J. Currie, and V. Duque (2018). Childhood circumstances and adult outcomes: Act ii. *Journal of Economic Literature* 56(4), 1360–1446.
- Anderson, M. L. (2008). Multiple inference and gender differences in the effects of early intervention: A reevaluation of the abecedarian, perry preschool, and early training projects. *Journal of the American Statistical Association* 103(484).
- Angrist, J. D. and J.-S. Pischke (2008). *Mostly Harmless Econometrics: An Empiricist’s Companion*. Princeton University Press.
- Attanasio, O., S. Cattan, E. Fitzsimons, C. Meghir, and M. Rubio-Codina (2015, February). Estimating the production function for human capital: Results from a randomized control trial in colombia. Working Paper 20965, National Bureau of Economic Research.
- Attanasio, O. P., C. Fernández, E. O. Fitzsimons, S. M. Grantham-McGregor, C. Meghir, and M. Rubio-Codina (2014). Using the infrastructure of a conditional cash transfer program to deliver a scalable integrated early child development program in colombia: cluster randomized controlled trial. *BMJ* 349, g5785.
- Baranov, V., S. Bhalotra, P. Biroli, and J. Maselko (2019). Maternal depression, women’s empowerment, and parental investment: Evidence from a randomized control trial. *American Economic Review*.
- Belfield, C. R., M. Nores, S. Barnett, and L. Schweinhart (2006). The high/scope perry preschool program cost–benefit analysis using data from the age-40 followup. *Journal of Human Resources* 41(1), 162–190.

- Bhalotra, S., M. Karlsson, and T. Nilsson (2017). Infant health and longevity: Evidence from a historical intervention in sweden. *Journal of the European Economic Association* 15(5), 1101–1157.
- Biroli, P., T. Boneva, A. Raja, and C. Rauh (2018). Parental beliefs about returns to child health investments.
- Boneva, T. and C. Rauh (2018). Parental beliefs about returns to educational investments—the later the better? *Journal of the European Economic Association* 16(6), 1669–1711.
- Bütikofer, A., K. V. Løken, and K. G. Salvanes (2019). Infant health care and long-term outcomes. *Review of Economics and Statistics* 101(2), 341–354.
- Butikofer, A., J. Riise, and M. Skira (2018). The impact of paid maternity leave on maternal health. *NHH Dept. of Economics Discussion Paper* (04).
- Campbell, F., G. Conti, J. J. Heckman, S. H. Moon, R. Pinto, E. Pungello, and Y. Pan (2014). Early childhood investments substantially boost adult health. *Science* 343(6178), 1478–1485.
- Carneiro, P. and R. Ginja (2014). Long-term impacts of compensatory preschool on health and behavior: Evidence from head start. *American Economic Journal: Economic Policy* 6(4), 135–73.
- Cascio, E. U. (2009). Do investments in universal early education pay off? long-term effects of introducing kindergartens into public schools. Working Paper 14951, National Bureau of Economic Research.
- Cascio, E. U. (2015). The promises and pitfalls of universal early education. *IZA World of Labor*.

- Conti, G., J. J. Heckman, and R. Pinto (2016). The effects of two influential early childhood interventions on health and healthy behaviour. *The Economic Journal* 126(596), F28–F65.
- Cooper, P. J. and L. Murray (1998). Postnatal depression. *BMJ* 316(7148), 1884–1886.
- Cunha, F., I. Elo, and J. Culhane (2013, June). Eliciting maternal expectations about the technology of cognitive skill formation. Working Paper 19144, National Bureau of Economic Research.
- Currie, J. and D. Thomas (1995). Does head start make a difference? *American Economic Review* 85(3), 341–364.
- De Haan, M. and E. Leuven (2016). Head start and the distribution of long term education and labor market outcomes. Technical report, CESifo Working Paper No 5870.
- Deming, D. (2009). Early childhood intervention and life-cycle skill development: Evidence from head start. *American Economic Journal: Applied Economics*, 111–134.
- Doyle, O., N. Fitzpatrick, J. Lovett, and C. Rawdon (2015). Early intervention and child physical health: Evidence from a dublin-based randomized controlled trial. *Economics & Human Biology* 19, 224 – 245.
- Due, J. J. and J. S. Madsen (2008). *Forligsmagere og Forumshoppere – Analyse af OK 2008 i Den Offentlige Sektor*. Jurist- og Økonomforbundets Forlag.
- Garces, E., D. Thomas, and J. Currie (2002). Longer-term effects of head start. *The American Economic Review* 92(4), 999–1012.
- García, J. L., J. J. Heckman, D. E. Leaf, and M. J. Prados (2016). The life-cycle benefits of an influential early childhood program. Working Paper 22993, National Bureau of Economic Research.

- García, J. L., J. J. Heckman, and A. L. Ziff (2017, May). Gender differences in the benefits of an influential early childhood program. Working Paper 23412, National Bureau of Economic Research.
- Gertler, P., J. Heckman, R. Pinto, A. Zanolini, C. Vermeersch, S. Walker, S. M. Chang, and S. Grantham-McGregor (2014). Labor market returns to an early childhood stimulation intervention in jamaica. *Science* 344(6187), 998–1001.
- Heckman, J., S. H. Moon, R. Pinto, P. Savelyev, and A. Yavitz (2010a). Analyzing social experiments as implemented: A reexamination of the evidence from the highscope perry preschool program. *Quantitative Economics* 1(1), 1–46.
- Heckman, J., R. Pinto, and P. Savelyev (2013). Understanding the mechanisms through which an influential early childhood program boosted adult outcomes. *American Economic Review* 103(6), 2052–86.
- Heckman, J. J., S. H. Moon, R. Pinto, P. A. Savelyev, and A. Yavitz (2010b). The rate of return to the highscope perry preschool program. *Journal of Public Economics* 94(1), 114–128.
- Hjort, J., M. Sølvssten, and M. Wüst (2017). Universal investment in infants and long-run health: Evidence from denmark’s 1937 home visiting program. *American Economic Journal: Applied Economics* 9(4), 78–104.
- Ibsen, C. L., T. P. Larsen, J. S. Madsen, and J. Due (2011). Challenging scandinavian employment relations: the effects of new public management reforms. *The International Journal of Human Resource Management* 22(11), 2295–2310.
- Kronborg, H., H. H. Sievertsen, and M. Wüst (2016). Care around birth, infant and mother health and maternal health investments – evidence from a nurse strike. *Social Science and Medicine* 150, 201 – 211.

- Lovejoy, M. C., P. A. Graczyk, E. O'Hare, and G. Neuman (2000). Maternal depression and parenting behavior: A meta-analytic review. *Clinical Psychology Review* 20(5), 561–592.
- Ludwig, J. and D. Miller (2007). Does head start improve children's life chances? evidence from a regression discontinuity design*. *The Quarterly Journal of economics* 122(1), 159–208.
- Masse, L. N. and W. S. Barnett (2002). A benefit-cost analysis of the abecedarian early childhood intervention. *Cost-Effectiveness and Educational Policy, Larchmont, NY: Eye on Education, Inc*, 157–173.
- Olds, D. L., C. R. Henderson, R. Chamberlin, and R. Tatelbaum (1986). Preventing child abuse and neglect: A randomized trial of nurse home visitation. *Pediatrics* 78(1), 65–78.
- Olds, D. L., C. R. Henderson, R. Cole, J. Eckenrode, D. Kitzmann, Harriet Luckey, L. Pettitt, K. Sidora, P. Morris, and J. Powers (1998). Long-term effects of nurse home visitation on childrens criminal and antisocial behavior: 15-year follow-up of a randomized controlled trial. *JAMA* 280(14), 1238–1244.
- Olds, D. L., J. Robinson, R. O'Brien, D. W. Luckey, L. M. Pettitt, C. R. Henderson, R. K. Ng, K. L. Sheff, J. Korfmacher, S. Hiatt, and A. Talmi (2002). Home visiting by paraprofessionals and by nurses: A randomized, controlled trial. *Pediatrics* 110(3), 486–496.
- Paulson, J. F., S. Dauber, and J. A. Leiferman (2006). Individual and combined effects of postpartum depression in mothers and fathers on parenting behavior. *Pediatrics* 118(2), 659–668.
- Persson, P. and M. Rossin-Slater (2019). When dad can stay home: Fathers' workplace flexibility and maternal health. Technical report, National Bureau of Economic Research.
- Sandner, M. (2019). Effects of early childhood intervention on fertility and maternal employment: Evidence from a randomized controlled trial. *Journal of Health Economics* 63, 159–181.

- Sandner, M., T. Cornelissen, T. Jungmann, and P. Herrmann (2018). Evaluating the effects of a targeted home visiting program on maternal and child health outcomes. *Journal of Health Economics* 58, 269 – 283.
- Schweinhart, L. J., J. Montie, Z. Xiang, W. S. Barnett, C. R. Belfield, and M. Nores (2005). *Lifetime effects: The High/Scope Perry Preschool study through age 40*. Ypsilanti, MI: High/Scope Press.
- Sundhedsstyrelsen (2007). Primary preventive care for children and youth - national guidelines [forebyggende sundhedsordninger for børn og unge - retningslinier]. Technical report, The Danish National Board of Health.
- The Danish National Board of Health [Sundhedsstyrelsen] (2018). The child vaccination program - yearly report 2017 [børnevaccinationsprogrammet - Årsrapport 2017]. Technical report, The Danish National Board of Health.
- Thompson, O. (2017). Head start’s long-run impact: Evidence from the program’s introduction. *Journal of Human Resources*, 0216–7735r1.
- Vaithianathan, R., M. Wilson, T. Maloney, and S. Baird (2016). *The Impact of the Family Start Home Visiting Programme on Outcomes for Mothers and Children: A Quasi-Experimental Study*. Ministry of Social Development.
- Wachs, T. D., M. M. Black, and P. L. Engle (2009). Maternal depression: a global threat to children’s health, development, and behavior and to human rights. *Child Development Perspectives* 3(1), 51–59.
- Wüst, M. (2012). Early interventions and infant health: Evidence from the danish home visiting program. *Labour Economics* 19, 484–495.

A Appendix - For online publication





Fig. A1 Average number of universal and total nurse home visits for children in the treated and control period


Notes: Average number of visits is calculated for children in (September 18, 2007 - April 15, 2008) and control period (September 17, 2008 - April 15, 2009) in 21 equally sized 10-day bins.




(a) Mothers GP contacts, 365 days prior birth (b) Mother psychiatric diagnosis, prior to birth

Fig. A2 Common trend in pre-treatment health outcome: Number of mothers' GP contacts (a) and indicator if mother received a psychiatric diagnosis (b) in the year prior to birth


Notes: See notes to Figure 3. Treated period: September 18, 2007 - April 15, 2008. Control period: September 17, 2008 and 2009 - April 15, 2009 and 2010).




(a) Midwife visits



(b) Prenatal GP contacts



(c) Days admitted at birth



(d) C-sections

Fig. A3 Care around birth for the treated period and control periods

Notes: Panel (a) shows the number of prenatal midwife contacts, panel (b) shows the number of prenatal GP consultations, panel (c) shows the number of days admitted to hospital at birth and panel (d) shows the C-section rate. See notes to Figure 3. The sample includes children who were born in Copenhagen in the treated period (September 18, 2007 - April 15, 2008) and control periods (September 17, 2008 and 2009 - April 15, 2009 and 2010).




Fig. A4 Density of births

Notes: The figure show the density of births for 20 equally sized bins and a window 258 days prior to the beginning of the strike and 60 after the beginning of the strike. Grey bars are the strike exposed period and bars with black outline are children born on same dates the two following years. The vertical lines indicate the data period of our main analyses (treated period: September 18, 2007 - April 15, 2008 and control periods: September 17, 2008 and 2009 - April 15, 2009 and 2010).






Fig. A5 Share of children observed as Copenhagen residents on January 1 in the treated and control periods


Notes: See notes to Figure 3. The sample includes children who were born in Copenhagen in the treated period (September 18, 2007 - April 15, 2008) and control periods (September 17, 2008 and 2009 - April 15, 2009 and 2010).




(a) Total GP contacts - Not Health educated



(b) Total GP contacts - Health educated



(c) Emergency GP contacts - Not health educated



(d) Emergency GP contacts - Health educated

Fig. A6 Accumulated GP contacts at age four for children born in the treated period (September 18, 2007 - April 15, 2008) and the control periods (September 17, 2008 and 2009 - April 15, 2009 and 2010) and whether parents are educated in health

Notes: The figure shows the relationship between date of birth and accumulated GP contacts at age four by parental background. See Figure 2 for further details.

Table A1 Nurse home visiting in the municipality of Copenhagen

Visit (and eligibility)	Timing
Universal visits	
Initial visit (A)	0-14 days after birth
2-months visit (B)	After two months of life
4-months visit (C)	After four months of life
8-months visit (D)	After eight months of life
Visits on parental demand	
Pregnancy visit	30th week of gestation
Maternity visit	Immediately after birth. Home births and early discharge
1,5-year visit	1,5 years after birth
3-year visit	3 years after birth
Targeted offer (at-risk families)	
Extra home visits	At discretion of nurses

Notes: Source: Official guidelines for the Copenhagen NHV program.

Table A2 Balance testing: Parental covariates as outcome

Days	Prim. school, mother (1)	Prim. school, father (2)	Income, mother (3)	Income, father (4)	Cohabiting (5)	Married (6)	Young mother (7)	Young father (8)
180-151	-0.012 (0.024)	-0.023 (0.024)	-8.567 (10.197)	-123.961 (140.780)	0.034 (0.031)	-0.028 (0.032)	0.013 (0.011)	0.008 (0.007)
150-121	-0.021 (0.024)	-0.001 (0.025)	-1.703 (9.999)	-137.039 (140.867)	-0.008 (0.031)	-0.025 (0.032)	0.004 (0.010)	0.004 (0.007)
120-91	0.008 (0.025)	-0.039* (0.023)	10.751 (10.408)	-113.671 (141.624)	0.045 (0.031)	-0.015 (0.033)	-0.012 (0.011)	-0.002 (0.006)
90-61	0.021 (0.025)	0.007 (0.024)	-1.872 (10.817)	-115.782 (140.864)	0.046 (0.029)	0.017 (0.034)	0.014 (0.011)	0.007 (0.006)
60-31	-0.034 (0.024)	-0.010 (0.024)	-2.525 (10.205)	-107.583 (140.496)	0.050* (0.029)	-0.029 (0.032)	0.011 (0.010)	0.008 (0.006)
30-1	-0.014 (0.024)	-0.034 (0.023)	11.237 (28.824)	-86.723 (140.922)	0.034 (0.029)	-0.015 (0.033)	0.015 (0.011)	-0.003 (0.006)
Obs.	12568	12568	12568	12568	12568	12568	12568	12332

Notes: Each column shows the estimates from separate regressions. The coefficients are for the interactions of 30-day bins and a strike indicator. All regressions include period and bin fixed effects. The sample includes children who were born in Copenhagen in the treated period (September 18, 2007 - April 15, 2008) and in control periods (September 17, 2008 and 2009 - April 15, 2009 and 2010). Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A3 Balance testing: Covariates at birth as outcome

Days	Hosp. nights at birth (1)	Midwife contacts (2)	C- section (3)	Home birth (4)	Preterm birth (5)	Low birth weight (6)	Head size (7)	Female child (8)
180-151	-0.912 (0.640)	-0.018 (0.110)	-0.009 (0.030)	0.000 (0.002)	-0.003 (0.018)	-0.019 (0.017)	0.088 (0.129)	0.025 (0.035)
150-121	-0.308 (0.636)	0.105 (0.108)	-0.005 (0.029)	0.003 (0.004)	-0.034** (0.017)	-0.019 (0.016)	0.018 (0.137)	0.060* (0.035)
120-91	-0.716 (0.716)	0.023 (0.116)	-0.003 (0.030)	-0.002 (0.002)	-0.033* (0.018)	-0.040** (0.016)	-0.070 (0.129)	0.038 (0.036)
90-61	-0.703 (0.624)	-0.004 (0.118)	0.010 (0.030)	-0.000 (0.004)	-0.023 (0.017)	-0.021 (0.015)	-0.039 (0.123)	0.047 (0.036)
60-31	-0.675 (0.644)	0.090 (0.116)	-0.021 (0.028)	0.001 (0.003)	-0.019 (0.017)	-0.011 (0.016)	0.027 (0.127)	0.071** (0.035)
30-1	-0.627 (0.638)	-0.083 (0.103)	-0.001 (0.029)	-0.003 (0.003)	-0.037** (0.016)	-0.022 (0.015)	0.249* (0.137)	0.058 (0.035)
Obs.	12537	12409	12568	12568	12518	12515	12332	12568

Notes: See notes for Table A2. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A4 Variable means, population of children born in Copenhagen and Denmark.

	Denmark Excl. CPH		CPH	
	Mean	Obs.	Mean	Obs.
Cohabitation	0.86	115578	0.78	17949
Married	0.47	115302	0.39	17917
Prim. school, mother	0.18	111553	0.13	17054
Uni. degree, mother	0.13	111553	0.33	17054
Student, mother	0.03	114562	0.05	17927
Employed, mother	0.81	114562	0.79	17927
Prim. school, father	0.19	110697	0.15	16561
Uni. degree, father	0.13	110697	0.33	16561
Student, father	0.01	113425	0.03	17334
Employed, father	0.90	113425	0.86	17334
Danish, mother	0.86	116827	0.76	18302
Danish, father	0.87	115578	0.75	17949
Young mother	0.05	116827	0.02	18302
Young father	0.02	115578	0.01	17949
Income, mother	255.79	114550	267.55	17926
Income, father	367.66	112391	361.10	17179
Length child	51.72	113575	51.66	17849
Low birth weight	0.05	114518	0.05	18021
Preterm birth	0.07	114637	0.06	18020
Head size	34.94	112024	34.79	17746
First time mothers	0.43	112743	0.62	17967
Multiple birth	0.04	116827	0.04	18302
C-section	0.22	116827	0.22	18302
No. of hospital nights at birth, child	3.83	114819	3.83	18070
Home birth	0.01	116827	0.01	18302
Midwife visits	4.80	111599	4.76	17814
Smoking status, Mother	0.17	114653	0.09	18020
BMI mom	24.46	107368	22.92	17424
Height mom	167.98	108542	167.88	17557

Notes: The Copenhagen sample includes all children born in Copenhagen in the periods: September 18, 2007 - April 15, 2008 and September 17, 2008 and 2009 - April 15, 2009 and 2010. The Denmark samples includes all children born in the same periods in Denmark, excluding Copenhagen.

Table A5 Additional child health outcomes: Effects of strike exposure on child hospitalization and outpatient contacts

Days	(1)	(2)	(3)	(4)	(5)	(6)
	Hospital adm. 1st year	Hospital adm. 2nd year	Hospital adm. 3-4 years	Outpat. cont. 1st year	Outpat. cont. 2nd year	Outpat. cont. 3-4 years
180-151	0.008 (0.030)	0.007 (0.027)	0.032 (0.028)	0.009 (0.035)	-0.029 (0.033)	0.024 (0.036)
150-121	-0.032 (0.030)	-0.009 (0.028)	-0.015 (0.028)	0.038 (0.035)	-0.026 (0.032)	0.035 (0.036)
120-91	0.001 (0.031)	0.015 (0.028)	-0.032 (0.028)	-0.012 (0.035)	-0.021 (0.033)	-0.051 (0.036)
90-61	0.037 (0.031)	0.024 (0.029)	-0.015 (0.028)	-0.003 (0.035)	-0.032 (0.033)	-0.029 (0.036)
60-31	0.004 (0.030)	0.067** (0.028)	0.013 (0.028)	-0.027 (0.035)	-0.005 (0.032)	0.000 (0.035)
30-1	0.036 (0.030)	0.078*** (0.029)	0.007 (0.028)	-0.060* (0.034)	0.025 (0.033)	-0.002 (0.036)
Control group mean	0.27	0.19	0.18	0.39	0.27	0.41
Obs.	12078	11982	11709	12078	11982	11709

Notes: See notes to table 3. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A6 Heterogeneity: Effects of strike exposure on total GP contacts by parental education and parity

Days	Not health educ.		Health educ.		Higher parity		First-borns	
	Total GP 1st year (1)	Total GP 2-4 years (2)	Total GP 1st year (3)	Total GP 2-4 years (4)	Total GP 1st year (5)	Total GP 2-4 years (6)	Total GP 1st year (7)	Total GP 2-4 years (8)
180-151	0.478 (0.569)	0.737 (1.081)	0.723 (1.241)	2.362 (2.333)	0.564 (0.745)	-1.207 (1.489)	0.478 (0.700)	2.134 (1.317)
150-121	0.153 (0.554)	0.921 (1.046)	1.313 (1.477)	1.096 (2.859)	0.198 (0.801)	-1.147 (1.490)	0.381 (0.679)	2.225* (1.307)
120-91	-0.078 (0.570)	-0.389 (1.015)	-1.318 (1.358)	0.543 (2.522)	-0.482 (0.807)	-0.710 (1.498)	0.020 (0.693)	0.509 (1.236)
90-61	0.754 (0.601)	1.594 (1.040)	0.459 (1.387)	0.318 (2.619)	0.520 (0.754)	0.462 (1.433)	0.842 (0.766)	2.002 (1.307)
60-31	0.262 (0.579)	2.347** (1.051)	0.517 (1.267)	2.244 (2.410)	0.339 (0.774)	1.304 (1.415)	0.397 (0.716)	3.214** (1.318)
30-1	1.924*** (0.611)	3.484*** (1.101)	1.375 (1.326)	1.234 (2.159)	2.319*** (0.831)	1.574 (1.462)	1.511** (0.738)	3.873*** (1.341)
Control group mean	4.72	20.95	3.52	18.75	4.07	17.88	4.91	22.59
Observations	10445	10114	1633	1584	4750	4605	7328	7093

Notes: See notes for Table 3. Column labels indicate the relevant subgroup and outcome variable studied. Columns (1)-(4) split the sample by parental educational background in a health-related field (either one of the parents are educated as a doctor, midwife, nurse or pedagogue). Columns (5)-(8) split the sample by parity of the child. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A7 Heterogeneity: Effects of strike exposure on total GP contacts, interacted model

	Health education		Parity	
	Total GP 1st year (1)	Total GP 2-4 years (2)	Total GP 1st year (3)	Total GP 2-4 years (4)
Days				
180-151	-0.153 (1.336)	1.228 (2.657)	-0.022 (1.018)	3.279* (1.987)
150-121	0.953 (1.541)	0.473 (3.032)	0.160 (1.051)	3.289* (1.980)
120-91	-1.052 (1.468)	1.807 (2.754)	0.600 (1.063)	1.093 (1.931)
90-61	-0.173 (1.493)	-0.165 (2.797)	0.177 (1.074)	1.282 (1.932)
60-31	0.168 (1.355)	0.852 (2.633)	0.101 (1.049)	1.817 (1.927)
30-1	-0.773 (1.454)	-2.437 (2.425)	-0.757 (1.111)	2.137 (1.982)
Observations	12078	11698	12078	11698

Notes: Each column shows the estimates from separate regressions. Column labels indicate the relevant subgroup of our sample. The coefficients are for the interactions of 30 day bins, a strike indicator and subgroup. All regressions include period, bin fixed effects and the interaction between bin indicators and strike exposure and full interactions between those and subgroup indicator. Regressions also include all control variables (see notes for Table 3). The sample includes children who were born in Copenhagen in the treated period (September 18, 2007 - April 15, 2008) and in control periods (September 17, 2008 and 2009 - April 15, 2009 and 2010). Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A8 Heterogeneity: Effects of strike exposure on total GP contacts at age four

	Gender		Initial health		SES		Smoking, mother	
	Boys (1)	Girls (2)	Not poor (3)	poor (4)	High (5)	Low (6)	No (7)	Yes (8)
Days								
180-151	3.288* (1.802)	-0.946 (1.846)	1.094 (1.366)	5.329 (3.803)	2.289 (1.424)	-0.897 (2.733)	1.038 (1.326)	4.902 (5.182)
150-121	0.263 (1.753)	2.525 (1.870)	1.410 (1.333)	1.095 (4.672)	3.238** (1.442)	-3.478 (2.633)	1.211 (1.326)	3.083 (4.676)
120-91	0.596 (1.770)	-1.493 (1.749)	-0.919 (1.307)	8.509** (4.152)	1.232 (1.389)	-4.064 (2.615)	-0.892 (1.284)	4.944 (4.908)
90-61	3.727** (1.837)	0.616 (1.807)	2.263* (1.358)	1.520 (4.148)	3.125** (1.484)	-0.242 (2.569)	1.846 (1.360)	5.659 (4.350)
60-31	3.580* (1.918)	1.799 (1.742)	1.911 (1.330)	13.082** (5.538)	3.869*** (1.443)	0.267 (2.682)	2.224 (1.363)	8.323** (4.109)
30-1	6.457*** (1.871)	2.617 (1.857)	4.336*** (1.379)	8.088 (5.189)	5.001*** (1.447)	4.105 (2.914)	4.344*** (1.374)	8.438* (4.834)
Control group mean	26.35	24.00	25.17	26.97	24.68	26.51	25.05	27.99
Observations	6085	5644	10803	926	8381	3348	10681	1048

Notes: See notes to Table 3 and A6. Columns (1)-(2) split the sample by child gender. Columns (3)-(4) split the sample by initial health (low birth weight, premature birth or complications during birth). Columns (5)-(6) split the sample by parental socio-economic status (SES). A low SES background is a child born to parents with either incomes in the bottom decile, below age 21 at birth or with only primary schooling. Columns (7)-(8) split the sample by whether the mother smoked during pregnancy. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A9 Parental investments: Effects of strike exposure on participation in preventive health checks

	(1)	(2)	(3)	(4)	(5)	(6)
	Prev. care, 5 weeks	Prev. care, 5 months	Prev. care, 12 months	Prev. care, 2 years	Prev. care, 3 years	Prev. care, 4 years
Days						
180-151	0.002 (0.022)	0.005 (0.019)	0.005 (0.018)	0.065* (0.034)	0.057 (0.035)	0.034 (0.030)
150-121	0.007 (0.021)	-0.008 (0.018)	0.011 (0.019)	0.043 (0.034)	0.036 (0.035)	0.005 (0.031)
120-91	-0.009 (0.022)	-0.008 (0.019)	-0.009 (0.019)	0.010 (0.035)	-0.034 (0.036)	-0.017 (0.031)
90-61	0.015 (0.021)	0.004 (0.020)	0.012 (0.018)	0.107*** (0.034)	0.099*** (0.036)	0.039 (0.031)
60-31	0.017 (0.021)	-0.014 (0.019)	0.029* (0.018)	0.034 (0.033)	0.090*** (0.034)	0.018 (0.030)
30-1	0.012 (0.020)	-0.000 (0.019)	0.017 (0.018)	0.056 (0.034)	0.083** (0.035)	0.037 (0.030)
Control	0.92	0.93	0.93	0.66	0.58	0.79
group						
mean						
Obs.	12078	12078	12078	11982	11832	11729

Notes: See notes for Table 3. Outcomes are indicators for participation in each consultation in the preventive health care program. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A10 Parental investments: Effects of strike exposure on participation in the infant vaccination program

	(1) Vacc., 1st round	(2) Vacc., 2nd round	(3) Vacc., 3rd round
Days			
180-151	-0.025 (0.025)	-0.015 (0.023)	-0.036* (0.022)
150-121	-0.005 (0.024)	-0.032 (0.023)	-0.039* (0.022)
120-91	0.013 (0.024)	-0.009 (0.023)	-0.045** (0.023)
90-61	-0.011 (0.025)	-0.010 (0.024)	-0.021 (0.023)
60-31	-0.018 (0.024)	-0.026 (0.023)	0.017 (0.022)
30-1	0.006 (0.024)	0.001 (0.023)	-0.034 (0.022)
Control group mean	0.90	0.91	0.91
Obs.	12078	12078	12078

Notes: See notes for Table 3. Outcomes are indicators for participation in each vaccination round scheduled within the first year of a child's life. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A11 Robustness: Effects of strike exposure on child GP contacts without pre-treatment covariates

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Total GP	Total GP	Total GP	Total GP	Ordin. GP	Ordin. GP	Ordin. GP	Emerg. GP	Emerg. GP	Emerg. GP
1st year	2nd year	3-4 years	1st year	2nd year	3-4 years	1st year	2nd year	3-4 years	3-4 years
Days									
180-151	0.470 (0.513)	0.025 (0.513)	0.556 (0.595)	0.166 (0.344)	-0.211 (0.318)	0.359 (0.387)	0.304 (0.221)	0.236 (0.296)	0.197 (0.313)
150-121	0.312 (0.514)	0.305 (0.505)	0.505 (0.598)	0.174 (0.339)	0.142 (0.317)	0.428 (0.385)	0.138 (0.228)	0.163 (0.289)	0.077 (0.313)
120-91	-0.294 (0.518)	-0.334 (0.486)	-0.248 (0.563)	-0.272 (0.348)	-0.053 (0.305)	0.036 (0.378)	-0.022 (0.218)	-0.281 (0.283)	-0.284 (0.295)
90-61	0.808 (0.545)	0.763 (0.510)	1.054* (0.581)	0.370 (0.373)	0.469 (0.327)	0.687* (0.387)	0.438** (0.223)	0.294 (0.284)	0.366 (0.305)
60-31	0.268 (0.520)	1.571*** (0.501)	0.400 (0.573)	0.068 (0.346)	0.896*** (0.317)	0.257 (0.376)	0.201 (0.222)	0.675** (0.288)	0.143 (0.300)
30-1	1.750*** (0.550)	1.362*** (0.516)	1.033* (0.584)	1.077*** (0.371)	0.781** (0.328)	0.585 (0.390)	0.674*** (0.233)	0.581** (0.288)	0.448 (0.297)
Control	4.55	10.35	10.22	3.09	6.89	7.04	1.47	3.46	3.18
group									
mean									
Obs.	12568	12464	12177	12568	12464	12177	12568	12464	12177

Notes: See notes to Table 3. We estimate the effects of strike exposure without pre-treatment covariates. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A12 Robustness: Effects of strike exposure on child GP contacts, larger bin size - 35 days

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
	Total GP 1st year	Total GP 2nd year	Total GP 3-4 years	Ordin. GP 1st year	Ordin. GP 2nd year	Ordin. GP 3-4 years	Emerg. GP 1st year	Emerg. GP 2nd year	Emerg. GP 3-4 years
Days									
175-141	0.266 (0.476)	0.318 (0.472)	0.529 (0.557)	0.107 (0.318)	0.089 (0.294)	0.336 (0.363)	0.159 (0.209)	0.229 (0.271)	0.193 (0.291)
140-106	-0.544 (0.483)	-0.334 (0.472)	0.189 (0.540)	-0.377 (0.323)	-0.027 (0.296)	0.277 (0.357)	-0.168 (0.207)	-0.307 (0.270)	-0.088 (0.280)
105-71	0.174 (0.508)	0.412 (0.463)	0.206 (0.524)	0.041 (0.347)	0.355 (0.293)	0.119 (0.354)	0.133 (0.211)	0.057 (0.265)	0.087 (0.274)
70-36	0.411 (0.490)	1.450*** (0.465)	0.805 (0.523)	0.141 (0.328)	0.788*** (0.298)	0.419 (0.348)	0.270 (0.205)	0.662** (0.265)	0.386 (0.271)
35-1	1.454*** (0.512)	1.579*** (0.485)	0.977* (0.536)	0.875** (0.346)	1.052*** (0.308)	0.552 (0.363)	0.579*** (0.218)	0.527* (0.272)	0.425 (0.270)
Control group mean	4.55	10.35	10.22	3.09	6.89	7.04	1.47	3.46	3.18
Observations	2078	11982	11709	12078	11982	11709	12078	11982	11709

Notes: See notes to Table 3. We increase the bin size to 35 days. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A13 Robustness: Effects of strike exposure on child GP contacts, smaller bin size - 21 days

Days	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
	Total GP 1st year	Total GP 2nd year	Total GP 3-4 years	Ordin. GP 1st year	Ordin. GP 2nd year	Ordin. GP 3-4 years	Emerg. GP 1st year	Emerg. GP 2nd year	Emerg. GP 3-4 years
189-169	0.886 (0.604)	0.00730 (0.599)	1.000 (0.683)	0.463 (0.407)	-0.0659 (0.381)	0.534 (0.453)	0.423 (0.258)	0.0732 (0.340)	0.465 (0.356)
168-148	0.736 (0.605)	0.373 (0.615)	0.442 (0.722)	0.501 (0.411)	0.0792 (0.378)	0.417 (0.470)	0.235 (0.258)	0.294 (0.353)	0.0249 (0.379)
147-127	0.293 (0.622)	0.491 (0.609)	0.324 (0.717)	0.297 (0.409)	0.437 (0.384)	0.422 (0.474)	-0.00421 (0.277)	0.0540 (0.346)	-0.0976 (0.362)
126-106	-0.134 (0.613)	-0.409 (0.600)	0.584 (0.691)	-0.146 (0.419)	-0.0567 (0.381)	0.500 (0.461)	0.0122 (0.249)	-0.353 (0.337)	0.0844 (0.368)
105-85	0.657 (0.650)	0.731 (0.587)	0.297 (0.670)	0.383 (0.440)	0.504 (0.377)	0.198 (0.454)	0.274 (0.279)	0.227 (0.338)	0.0991 (0.353)
84-64	0.693 (0.643)	0.260 (0.606)	0.583 (0.676)	0.344 (0.444)	0.259 (0.387)	0.400 (0.461)	0.349 (0.256)	0.00154 (0.333)	0.183 (0.347)
63-43	1.082* (0.627)	1.740*** (0.595)	0.965 (0.675)	0.629 (0.423)	1.053*** (0.388)	0.607 (0.454)	0.453* (0.260)	0.687** (0.336)	0.358 (0.349)
42-22	0.527 (0.636)	1.759*** (0.620)	1.435** (0.697)	0.245 (0.426)	1.172*** (0.393)	0.869* (0.459)	0.282 (0.274)	0.588* (0.347)	0.566 (0.363)
21-1	2.638*** (0.665)	1.749*** (0.623)	1.163* (0.701)	1.687*** (0.453)	0.942** (0.396)	0.567 (0.481)	0.951*** (0.277)	0.807** (0.350)	0.596* (0.349)
Control group mean	4.55	10.35	10.22	3.09	6.89	7.04	1.47	3.46	3.18
Observations	12078	11982	11709	12078	11982	11709	12078	11982	11709

Notes: See notes to Table 3. We reduce the bin size to 21 days. Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A14 Placebo test: The effect of strike exposure on child health measured as accumulated GP contacts by type, data for the two control years 2009 and 2010

	(1)	(2)	(3)	(4)	(5)	(6)
	Total GP 1st year	Total GP 2-4 years	Ordin. GP 1st year	Ordin. GP 2-4 years	Emerg. GP 1st year	Emerg. GP 2-4 years
Days						
180-151	0.270 (0.621)	-0.505 (1.097)	-0.076 (0.411)	0.153 (0.696)	0.346 (0.267)	-0.658 (0.589)
150-121	-0.154 (0.623)	-1.188 (1.084)	-0.455 (0.399)	-0.968 (0.670)	0.300 (0.284)	-0.220 (0.599)
120-91	-0.990 (0.637)	0.315 (1.096)	-1.019** (0.414)	0.632 (0.691)	0.029 (0.282)	-0.317 (0.592)
90-61	-0.341 (0.629)	0.678 (1.089)	-0.387 (0.420)	0.651 (0.688)	0.047 (0.265)	0.027 (0.585)
60-31	0.032 (0.634)	-0.494 (1.083)	0.057 (0.415)	0.054 (0.694)	-0.025 (0.279)	-0.548 (0.583)
30-1	-0.991 (0.623)	-0.529 (1.069)	-0.863** (0.417)	0.114 (0.699)	-0.129 (0.263)	-0.643 (0.552)
Control group mean	4.55	20.65	3.09	13.98	1.47	6.67
Obs.	8203	7941	8203	7941	8203	7941

Notes: Each column shows the estimates from separate regressions. The coefficients are for the interactions of 30 day bins and a strike indicator. All regressions include period and bin fixed effects, as well as control variables (see notes for Table 3). The sample includes children who were born in Copenhagen in the placebo treated period (September 17, 2008 - April 15, 2009) and in control period (September 17, 2009 - April 15, 2010). Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.

Table A15 Effect of strike exposure on child and mother health measured as accumulated and yearly total GP fees, Euro

	(1) Total GP mother and child 1st year	(2) Total GP mother and child 2nd year	(3) Total GP mother and child 3-4 years	(4) Total GP mother and child <4y
Days				
180-151	8.586 (14.174)	9.741 (13.573)	17.876 (19.877)	33.850 (39.289)
150-121	10.476 (13.846)	29.762** (13.437)	30.077 (19.355)	66.267* (38.443)
120-91	0.230 (14.013)	15.548 (13.329)	1.902 (18.994)	12.530 (37.556)
90-61	8.850 (14.388)	40.319*** (13.619)	37.636* (19.205)	87.099** (38.453)
60-31	14.918 (14.229)	52.904*** (13.726)	31.088 (19.487)	94.165** (39.242)
30-1	46.565*** (14.761)	61.200*** (13.752)	48.826** (19.190)	154.281*** (39.327)
Control group mean	240.22	307.22	426.41	978.01
Obs.	12078	11982	11709	11698

Notes: See notes for Table 3. GP fees are measured in Euro (2015-prices). Robust standard errors in parentheses. *** $p < 0.01$, ** $p < 0.05$ and * $p < 0.10$.